

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) EP 1 006 183 A1

(12) EUROPEAN PATENT APPLICATION

(43) Date of publication:
07.06.2000 Bulletin 2000/23

(21) Application number: 98122969.3

(22) Date of filing: 03.12.1998

(51) Int. Cl.⁷: C12N 15/12, C07K 14/705,
C12N 1/21, C12N 15/70,
G01N 33/53, G01N 33/68,
A61K 38/17, C07K 17/00

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(71) Applicant:
Max-Planck-Gesellschaft zur Förderung
der Wissenschaften e.V.
80539 München (DE)

(72) Inventor:
The designation of the Inventor has not yet been
filed

(74) Representative:
Böhm, Brigitte, Dipl.-Chem. Dr. et al
Kopernikusstrasse 9
81679 Munich (DE)

(54) Recombinant soluble Fc receptors

(57) Recombinant soluble Fc receptors according to the present invention are characterized by the absence of transmembrane domains, signal peptides and glycosylation. Such Fc receptors can easily be obtained by expressing respective nucleic acids in prokaryotic host cells and renaturation of the obtained inclusion bodies, which procedure leads to a very homogenous and pure product.

The products can be used for diagnostic as well as pharmaceutical applications and also for the generation of crystal structure data. Such crystal structure data can be used for the modelling of artificial molecules.

A further embodiment comprises coupling the Fc receptors according to the invention to solid materials like chromatography materials that can be used to separate and/or enrich antibodies.

EP 1 006 183 A1

BEST AVAILABLE COPY

Description

[0001] The present invention relates to recombinant soluble Fc receptors (FcR), recombinant nucleic acids coding for such Fc receptors, host cells containing corresponding nucleic acids as well as a process for the determination of the amount of antibodies of a certain type contained in the blood, plasma or serum of a patient, a process for the determination of the immune status of patients with chronic diseases of the immune system and a process for the screening of substances in view of their ability to act as inhibitors of the recognition and binding of antibodies to the respective cellular receptors. Further, the present invention is concerned with pharmaceutical compositions containing the recombinant soluble FcRs, the use of a crystalline preparation of recombinant soluble FcRs for the generation of crystal structure data of Fc receptors as well as FcR inhibitors and pharmaceutical compositions containing such FcR inhibitors.

[0002] A still further subject of the present invention is a recombinant Fc receptor coupled to a solid phase, e.g. a chromatography carrier material. The use of such chromatography material, which is another subject of the present invention, lies in the absorption of immunoglobulins from a body fluid of patients or from culture supernatants of immunoglobulin producing cells.

[0003] Fc receptors (FcRs) play a key role in defending the human organism against infections. After pathogens have gained access to the blood circulation they are opsonized by immunoglobulins (Igs). The resulting immunocomplexes bind due to their multivalency with high avidity to FcR bearing cells leading to clustering of the FcRs, which triggers several effector functions (Metzger, H., 1992A). These include, depending on the expressed FcR type and associated proteins, endocytosis with subsequent neutralization of the pathogens and antigen presentation, antibody-dependent cellular cytotoxicity (ADCC), secretion of mediators or the regulation of antibody production (Fridman et al, 1992; van de Winkel and Capel, 1993).

[0004] Specific FcRs exist for all Ig classes, the ones for IgG being the most abundant with the widest diversity. Together with the high affinity receptor for IgE (FcεRIa), FcγRI (CD64), FcγRII (CD32) and FcγRIIIa (CD16) occur as type I transmembrane proteins or in soluble forms (sFcRs) but also a glycosylphosphatidylinositol anchored form of the FcγRIII (FcγRIIIb) exists. Furthermore, FcγRs occur in various isoforms (FcγRIa, b1, b2, c; FcγRIIIa 1-2, b1-3, c) and alleles (FcγRIIIa1-HR, -LR; FcγRIIIb-NA1, -NA2) (van de Winkel and Capel, 1993). In contrast to the overall homologous extracellular parts, the membrane spanning and the cytoplasmic domains differ. They may be deleted entirely or be of a size of 8 kDa. They may contain either a 26 amino acid immunoreceptor tyrosine-based activation motif (ITAM) as in FcγRIIIa or a respective 13 amino acid inhibitory motif (ITIM) in FcγRIIIb involved in signal transduction (Amigorena et al, 1992).

[0005] Judged by the conserved spacing of cysteins, the extracellular part of the FcRs consists of three (FcγRI, CD64) or two (FcεRI, FcγRII, CD32 and FcγRIII, CD16) Ig-like domains (10 kDa/domain) and therefore belongs to the immunoglobulin super family. These highly glycosylated receptors are homologues, and the overall identity in amino acid sequence among the FcγRs and FcεRIa exceeds 50% in their extracellular regions. Nevertheless, the affinity of FcRs to their ligands varies widely. The higher affinity of $\approx 10^8 \text{M}^{-1}$ of the FcγRI to Fc-fragment is assigned to its third domain, while the other FcγRs with two domains have an affinity to IgG varying between 10^5 and 10^7M^{-1} . The affinity of the two domain FcεRIa to IgE exceeds these values by far with a constant of 10^{10}M^{-1} (Metzger, H., 1992B).

FcγRs are expressed in a defined pattern on all immunological active cells. FcγRI is constitutively expressed on monocytes and macrophages and can be induced on neutrophils and eosinophils. The physiological role of FcγRI is still unknown as the expression on monocytes is not vital (Ceuppens et al, 1988). The GPI anchored form of FcγRIII (FcγRIIIb) is exclusively expressed on granulocytes. Due to its missing cytoplasmic part, the signal transduction into the cell occurs solely via other transmembrane proteins like complement receptor type 3 (CR3) that can at least associate with FcγRIIIb (Zhou et al, 1993; Poo et al, 1995). FcγRIIIa is mainly expressed on monocytes and macrophages but only in conjunction with associated proteins (e.g. α- or γ-chains). FcγRII is the receptor with the widest distribution on immunocompetent cells and is mainly involved in the endocytosis of immunocomplexes.

[0006] FcγRIIIa and FcγRIIIb differ in their extracellular region by only 7% of the amino acid residues. Nevertheless, both forms can be distinguished by their binding characteristics to human and mouse IgG subclasses (van de Winkel and Capel, 1993) and their differing affinity to human IgGs (Sondermann et al, 1998A). The situation is rendered even more complicated by the high responder/low responder (HR/LR) polymorphism of FcγRIIIa named after the ability of T cells from some individuals to respond to murine IgG1-induced mitogenesis (Tax et al, 1983). Later, it was found that the two exchanges in the amino acid sequence between the LR and the HR form modify the ability to bind human IgG2, which leads to the suggestion that at least one of them is involved in IgG binding (Hogarth et al, 1992).

[0007] In contrast to the beneficial role FcRs play in the healthy individual, they also transmit the stimulation of the immune system in allergies (FcεRIa) or autoimmune diseases. Moreover, some viruses employ FcγRs to get access to cells like HIV (Homsy et al, 1989) and Dengue (Littau et al, 1990) or slow down the immune response by blocking FcγRs as in the case of Ebola (Yang et al, 1998) and Measles (Ravanel et al, 1997).

[0008] Hence, the object underlying the present invention was to provide receptors which are easy to produce and can advantageously be used for medical or diagnostic applications. Moreover, it was an object of the invention to pro-

vide soluble receptors exhibiting a binding specificity and activity which is analogous to that of the receptors occurring naturally in the human body and which, additionally, make it possible to produce crystals suitable for a structure determination.

[0009] This object is accomplished by recombinant soluble Fc receptors which consist only of the extracellular portion of the receptor and are not glycosylated. The receptors according to the present invention are therefore characterized by the absence of transmembrane domains, signal peptides and glycosylation.

[0010] Particularly preferred for the present invention are Fc γ or Fc ϵ receptors. This is because IgG and IgE molecules are characteristic for a multiplicity of diseases and conditions, so that their determination and possible ways of influencing them are of great interest. Figure 8 shows an alignment of amino acid sequences of the extracellular parts of some Fc γ R α s and Fc ϵ R α I. The FcRs according to the invention include all these sequences or parts thereof that still retain binding capacity to antibodies and/or proper crystallization.

[0011] In a particularly preferred embodiment of the invention the recombinant soluble FcR is a Fc γ R α IIb receptor. Further, it is particularly preferred that the receptor be of human origin. In a particularly preferred embodiment, it contains an amino acid sequence as shown in SEQ ID NO:1 or SEQ ID NO:2.

[0012] According to the present invention, the preparation of the soluble Fc receptors preferably takes place in prokaryotic cells. After such expression, insoluble inclusion bodies containing the recombinant protein form in prokaryotic cells, thus facilitating purification by separation of the inclusion bodies from other cell components before renaturation of the proteins contained therein takes place. The renaturation of the FcRs according to the present invention which are contained in the inclusion bodies can principally take place according to known methods. The advantage of the preparation in prokaryotic cells, the production of inclusion bodies and the thus obtained recombinant soluble Fc receptors make it possible to obtain a very pure and, in particular, also very homogeneous FcR preparation. Also because of the absence of glycosylation the obtained product is of great homogeneity.

[0013] Soluble Fc receptors hitherto produced by recombinant means particularly exhibited the disadvantage that a much more elaborate purification was required, since they were expressed in eukaryotic cells and, due to the glycosylation which is not always uniform in eukaryotic cells, these products were also less homogeneous.

[0014] The recombinant soluble Fc receptors according to the present invention even make it possible to produce crystals suitable for use in X-ray analysis, as shall be explained later on. The FcRs of the present invention moreover exhibit practically the same activity and specificity as the receptors naturally occurring in vivo.

[0015] A further subject matter of the present invention is a recombinant nucleic acid having a sequence coding for a recombinant soluble Fc receptor according to the present invention.

[0016] The nucleic acid according to the present invention may contain only the coding sequences or, additionally, vector sequences and, in particular, expression control sequences operatively linked to the sequence encoding the recombinant FcR, like promoters, operators and the like.

[0017] In a particularly preferred embodiment the nucleic acid of the present invention contains a sequence as shown in SEQ ID NO:3 or SEQ ID NO:4.

[0018] In these sequence protocols the atg start codons are in bold print and newly introduced restriction sites are underlined. For a comparison, SEQ ID NO:5 and SEQ ID NO:6 show the respective wild type sequences coding for Fc γ R α IIb and Fc ϵ R α Ia. SEQ ID NOs:7-9 show the wild type sequences for Fc γ R α I, Fc γ R α IIa and Fc γ R α III, which can be modified in a similar way as SEQ ID NO:3 and SEQ ID NO:4 to obtain FcRs according to the invention.

[0019] If the nucleic acid of the present invention contains vector sequences, then these are preferably sequences of one or several prokaryotic expression vectors, preferably of pET vectors. Any other known functions of expression vectors may also be contained in the recombinant nucleic acid according to the present invention if desired. These may, for instance, be resistance genes allowing for an effective selection of transformed host cells.

[0020] A still further subject matter of the present invention is a host cell containing a recombinant nucleic acid according to the present invention. As repeatedly mentioned above, the host cell preferably is a prokaryotic host cell, particularly an E. coli cell.

[0021] The recombinant soluble Fc receptors according to the present invention can be used for a multitude of examinations or applications because they specifically react with antibodies. In vivo, the soluble Fc receptors are powerful immunoregulators which, if present in elevated levels, result in a remarkable suppression of the immune system which leads to many partly known and partly not yet understood effects. Based on these effects, several applications of the Fc receptors according to the present invention are further subject matters of the present invention.

[0022] One such subject is a process for the determination of the amount of antibodies of a certain type in the blood or serum of a patient, which is characterized by the use of a recombinant soluble FcR according to the invention in an immunoassay, and the determination of the presence of FcR-antibody complexes. Such assay allows to screen for the presence of a certain kind of antibody and allows also for the determination of the amount of antibodies present in the blood, plasma or serum of a patient.

[0023] Any type of immunoassay is principally suitable for the use according to the present invention, as long as the presence of FcR-antibody complexes can thereby be detected. Both ELISA (enzyme-linked immunosorbent immu-

noassay), particularly sandwich assays, and RIA (radio-immunoassay) are suitable, but also competitive testing methods. In a preferred embodiment of the invention where the presence and/or the amount of IgE antibodies is to be examined, an Fc ϵ R is used as recombinant soluble receptor according to the present invention. In particular, this method is suited and advantageous for determining a predisposition or manifestation of an allergy.

5 [0024] Moreover, a method is preferred in which the presence of soluble FcRs is to be determined and, if required, quantified, an Fc γ R being used as recombinant soluble receptor according to the invention. By means of this test among others the immune status of patients with chronic diseases of the immune system can be determined in a competitive immunoassay. Chronic diseases in the sense of these processes are for instance AIDS, SLE (systemic lupus erythematosus), MM (multiple myeloma) or rheumatoid arthritis.

10 [0025] A further advantageous use of the receptor according to the present invention lies in the screening of substances in view of their ability to act as inhibitors of the recognition and binding of antibodies to the respective cellular receptors.

[0026] By means of modern screening techniques such as HTS (high throughput screening) in combination with multi-well microtiter plates and automatic pipetting apparatuses it is nowadays possible to simultaneously test a multi-
15 tude of substances for specific properties. As the FcRs according to the present invention can be easily produced at low cost, they can also be used in such series tests by which substances having an inhibiting effect can easily be identified.

[0027] Particularly preferred is such use according to which Fc receptors according to the present invention are used to find inhibitors capable of inhibiting the recognition and binding of the respective antibodies to the particular receptor of interest.

20 [0028] A further area of application of the substances according to the invention lies in the pharmaceutical field. Hence, a further subject matter of the invention is a pharmaceutical composition comprising as active agent a recombinant soluble FcR according to the invention. According to the present invention, this pharmaceutical composition may of course comprise conventional useful carrier and auxiliary substances. Such substances are known to the person of skill in the art, the mode of administration also having to be taken into account. The pharmaceutical composition of the
25 present invention can be advantageously used for the treatment or prevention of autoimmune diseases, allergies or tumor diseases.

[0029] Soluble forms of Fc receptors such as Fc γ RIII mediate isotype-specific regulation of B cell growth and immunoglobulin production. In a murine model of myeloma, sFcR suppresses growth and immunoglobulin production of tumor cells (Müller et al, 1985; Roman et al, 1988; Teillaud et al, 1990). Furthermore, sFcR binds to surface IgG on cultures of human IgG-secreting myeloma cells and effects suppression of tumor cell growth and IgG secretion. Prolonged
30 exposure of these cells to sFcR results in tumor cell cytotoxicity (Hoover et al, 1995).

[0030] Also, overreactions of the immune system in allergic reactions or due to massive antigen load might be reduced by, for example, intravenous application of soluble FcR (Ierino et al, 1993).

35 [0031] Therefore, a preferred pharmaceutical composition according to the invention for use in the treatment of AIDS, rheumatoid arthritis or multiple myeloma contains a recombinant soluble Fc γ receptor and, preferably, a receptor having the amino acid sequence as shown in SEQ ID NO:1.

[0032] It was also of great interest to obtain crystal structure data of Fc receptors. On the one hand, these are a key to the understanding of molecular mechanisms in immunocomplex recognition. On the other hand, these structural data can be used to find out common features in the structures of different Fc receptors and use the knowledge of the structures to generate inhibitors or identify and produce new artificial antibody receptors.
40

[0033] To obtain such crystal structure data, a crystalline preparation of the recombinant soluble Fc receptor according to the invention is used. The recombinant soluble FcRs according to the invention surprisingly can be obtained pure enough to produce crystals that give reliable X-ray structure determination data. Such crystallization was not possible with the hitherto produced receptor molecules, mostly due to their lack of homogeneity.

45 [0034] The stated applications are merely preferred embodiments of the use of the crystal structure data. Many other applications seem possible, too.

[0035] Suitably, the structural data for the generation and/or identification of inhibitors or new receptors, respectively, are used in a computer-aided modelling program. Software for computer-aided modelling is available to the man skilled in the art. That application is already described for structure identification or design of other substances.

50 [0036] Particularly preferred for the present invention are the structures as shown in the enclosed Examples and Figures for the respective receptors. Such structures can be used to design inhibitors, antagonists and artificial receptor molecules.

[0037] A still further subject matter of the present invention, therefore, is a FcR inhibitor which has a three-dimensional structure which is complementary to the recombinant soluble FcR according to the invention and inhibits the
55 binding of antibodies to FcRs.

[0038] What is important for the inhibitors of the invention is that, owing to their structure and specificity, they are capable of binding to the FcRs and thus prevent their normal binding to the constant parts of antibodies.

[0039] Preferably, such FcR inhibitors are small organic molecules which can easily be administered orally. They

might be an interesting alternative to cortisone in the treatment of autoimmune diseases and host/graft rejections. Such a molecule would also suppress reinfection rates with certain viruses, e.g. Dengue virus where the antibody coated virus is FcγRIIb dependent internalized (Littau et al, 1990), HIV where on CD4 positive T cells an antibody enhancement of HIV infection is mediated by FcγRIII (Hornsy et al, 1989), or Ebola where the virus secreted glycoprotein inhibits early neutrophil activation by blocking sFcγRIII which affects the host response to infection (Yang et al, 1998).

[0040] The development of inhibitors might also lead to substances that interfere with the recognition of IgE by their receptors. From the modelled structure of FcεRI, peptides have already been developed which inhibit mast cell degranulation in vitro. With the knowledge of the structures of the receptor or the receptor-antibody complex in atomic detail, a new possibility for a rational drug design is opened.

[0041] A further subject matter of the present invention therefore is a pharmaceutical composition containing as active agent an FcR inhibitor as mentioned above. Such pharmaceutical compositions may, for example, be used in the treatment or prevention of diseases which are due to overreactions or faulty reactions of the immune system, preferably the treatment or prevention of allergies, autoimmune diseases or anaphylactic shock.

[0042] A further subject of the present invention is the sFcR according to the invention, bound to a solid phase. Such heterogeneous receptors might be used for immunoassays or other applications where the receptor in an immobilized form can be used beneficially.

[0043] In a preferred embodiment of the invention the solid phase is a chromatography carrier material onto which the Fc receptor is fixed, e.g. sepharose, dextran sulfate etc. Such chromatography materials with Fc receptors bound thereto can beneficially be used for the adsorption of immunoglobulins from the blood, plasma or serum of patients or from the culture supernatant of immunoglobulin producing cells (meaning concentration, enrichment and purification of antibodies).

[0044] On the one hand, the antibodies bound to the chromatography material can be eluted and, for example, the immune status of a patient can thereby be determined. On the other hand, antibodies from the blood of a patient can thereby be enriched before carrying out further tests, which is a further preferred embodiment of the present invention. In many cases it is difficult to conduct diagnostic assays using blood samples if the latter contains only a very small number of the antibodies to be identified. By means of a concentration using a specific chromatographic column with Fc receptors according to the present invention, antibodies of interest can easily be concentrated and separated from many other substances which might disturb the test.

[0045] Basically, it is also possible to use a chromatography material according to the present invention in an extracorporeal perfusion system for lavage of the blood in case of certain diseases where the removal of antibodies plays a crucial role.

[0046] The following Examples are to further illustrate the invention in conjunction with the Figures.

Example 1

1.1 Cloning and Expression

[0047] The cDNA of human FcγRIIb2 (Engelhardt et al, 1990) was modified using mutagenous PCR (Dulau et al, 1989). Therefore, a forward primer was used for the introduction of a new start methionine after the cleavage site of the signal peptide within a *Nco*I site (5'-AAT AGA ATT CCA TGG GGA CAC CTG CAG CTC CC-3') while the reverse primer introduced a stop codon between the putative extracellular part and the transmembrane region followed by a *Sa*I site (5' CCC AGT GTC GAC AGC CTA AAT GAT CCC C-3'). The PCR product was digested with *Nco*I and *Sa*I, cloned into a pET11d expression vector (Novagen) and the proposed sequence was confirmed. The final construct was propagated in BL21 (DE3) (Grodberg and Dunn, 1988). For the overexpression of FcγRIIb a single colony of the transformed bacteria was inoculated in 5ml LB medium containing 100 µg ampicillin per ml (LB-Amp100) and incubated overnight at 37°C. The culture was diluted 200-fold in LB-Amp100 and incubation was continued until an OD600 of 0.7-0.9 was achieved. The overproduction of the protein was induced by adding IPTG to a final concentration of 1 mM. After a growing period of 4 hours the cells were harvested by centrifugation (30 mm, 4000 x g) and resuspended in sonification buffer (30 mM sodium phosphate, 300 mM sodium chloride, 0.02% sodium azide, pH 7.8). After addition of 0.1 mg lysozyme per ml suspension and incubation for 30 min at room temperature the sonification was performed on ice (Branson Sonifier, Danbury, CT; Macrotip, 90% output, 80% interval, 15 min). The suspension was centrifuged (30 min, 30,000 x g) and resuspended with a Dounce homogenizer in sonification buffer containing 0.5% LDAO. The centrifugation step and resuspension in LDAO containing buffer was repeated once before this procedure was repeated twice without LDAO. The purified inclusion bodies were stored at 4°C.

1.2 Refolding and purification of soluble human FcγRIIb (shFcγRIIb)

[0048] The purified inclusion bodies were dissolved to a protein concentration of 10 mg/ml in 6 M guanidine chlo-

ride, 100 mM 2-mercaptoethanol and separated from the insoluble matter by centrifugation. The refolding was achieved by rapid dilution. Therefore, one ml of the inclusion body solution was dropped under stirring within 15 hours into 400 ml of the refolding buffer (0.1 M TRIS/HCl, 1.4 M arginine, 150 mM sodium chloride, 5 mM GSH, 0.5 mM GSSG, 0.1 mM PMSF, 0.02% sodium azide, pH 8.5, 4°C). Afterwards, the mixture was stirred for 2-3 days until the concentration of free thiol groups was reduced to 1 mM by air oxidation as measured according to Ellman (Ellman, 1959). The solution was dialyzed against PBS and sterile filtered before it was concentrated 10-fold in a stirring cell equipped with a 3kD MWCO ultrafiltration membrane. The protein solution was applied to a hlgG sepharose column (50 mg hlgG per ml sepharose 4B). Unbound protein was washed out with 50 mM TRIS pH 8.0 before elution of FcγRIIb by pH jump (150mM sodium chloride, 100mM glycine, 0.02% sodium azide, pH 3.0). The eluate was immediately neutralized with 1 M TRIS pH 8.0. The FcγRIIb containing solution was concentrated and subjected to gel filtration on a Superdex-75 column equilibrated with crystallization buffer (2 mM MOPS 150 mM sodium chloride, 0.02% sodium azide pH 7.0). The fractions containing FcγRIIb were pooled, concentrated to 7 mg/ml and stored at -20°C.

1.3 Equilibrium gel filtration experiments

[0049] A Superdex75 column was connected to FPLC and equilibrated with PBS containing 10 μg shFcγRIIb per ml. Human Fc fragment was solved to a concentration of 1 μg/10 μl in the equilibration buffer and injected. The resulting chromatogram yielded a positive peak comprising the complex of the shFcγRIIb and the Fc fragment while the negative peak represents the lack of receptor consumed from the running buffer for complex formation.

1.4 Crystallization and data collection

[0050] Initial crystallization trials employing a 96 condition sparse matrix screen (Jancarik and Kim, 1991) were performed in sitting drops at 20°C using the vapor diffusion method. Occuring crystals were improved by changing the pH as well as the salt, precipitant and additive concentration. Diffraction data from suitable crystals was collected on an image plate system (MAR research) using graphite monochromated CuK_α radiation from a RU200b rotating anode generator (Rigaku) operated at 50 kV and 100 mA. The reflections were integrated with the program MOSFLM (Leslie, 1997) and subsequently the data was scaled, reduced and truncated to obtain the structure-factor amplitudes using routines from the CCP4 program suite (Collaborative Computational Project, 1994).

1.5 Summary of expression, purification and refolding of shFcγRIIb

[0051] The extracellular part of FcγRIIb was expressed in high levels under the control of a T7 promoter in the T7 RNA polymerase positive E. coli strain BL21/DE3 (Grodberg & Dunn, 1988). The protein was deposited in inclusion bodies, which were employed in the first purification step. The isolation of the inclusion bodies was started with an intense combined lysozyme/sonification procedure to open virtually all cells which would otherwise contaminate the product. The subsequent washing steps with the detergent LDAO, which has excellent properties in solving impurities but not the inclusion bodies itself already yielded a product with a purity of >90% (Fig. 1).

[0052] This product was used for refolding trials without further purification. The inclusion bodies were dissolved in high concentration of 2-mercaptoethanol and guanidine to ensure the shift of covalent and non-covalent aggregates to monomers. This solution was rapidly diluted with refolding buffer to minimize contacts between the unfolded protein molecules which would otherwise form aggregates. The use of arginine in the refolding buffer prevents the irreversible modification of side chains as often recognized with urea. After addition of the protein to the refolding buffer, the solution was stirred at 4°C until the concentration of free thiol groups was reduced to 1 mM, which was absolutely necessary as earlier dialysis resulted in an inactive product. In a second purification step the dialyzed and refolded FcγRIIb was bound to immobilized hlgG to remove minor fractions of E. coli proteins and inactive receptor. The protein was eluted with a pH jump and immediately neutralized. After this affinity chromatography step shFcγRIIb is essentially pure except for a minor contamination resulting from the coeluting IgG which leached out of the matrix even after repeated use (Fig. 1). The IgG as well as receptor multimers which are not visible in the reducing SDS-PAGE could easily be removed by gel filtration. Parallel to the removal of the contaminants in this step the buffer is quantitatively exchanged. This procedure ensures a defined composition of the protein solution as even slight variations can cause irreproducibility of the crystallization attempts or even inhibit the formation of crystals. Overall 6 mg pure protein could be gained per litre E. coli culture, which is about 10 % from the FcγRIIb content of the inclusion bodies.

[0053] N-terminal protein sequencing revealed the identity with the expected sequence H₂N-GTPAAP without detectable contamination. ESI-MS analysis showed that the final material used in crystallization trials is homogenous with respect to size. From the primary sequence the molecular weight was calculated to 20434 Da, which corresponds to 20429 Da found by mass spectroscopy. The discrepancy lies within the error of the instrument, and no additional peak for a species containing the leading methionine is found.

[0054] The crystallization of shFcγRIIb was performed in sitting drops using the vapor diffusion method. Initial trials with a sparse matrix screen (Jancarik & Kim, 1991) resulted already in small crystalline needles. Subsequent optimization of the preliminary crystallization condition by varying precipitant, salt, their concentration and pH led to the isolation of three different crystal forms. Orthorhombic crystals grew from mixture of 1.5 μl reservoir solution (33% PEG2000, 0.2 M sodium acetate, pH 5.4) with 3 μl of the protein solution. They appeared within 3 days and reached their final size of approximately 80 μm × 80 μm × 500 μm after one week. These crystals diffracted to 1.7 Å. Crystals could also be grown in two other space groups from reservoir solution containing 26% PEG8000, 0.2 M sodium acetate, pH 5.6, 5 mM Zn(OAc)₂, 100 mM sodium chloride (hexagonal form) and 26% PEG8000, 0.2 M NaOAc, pH 5.6, 10% (v/v) 1,4-Dioxan, 100 mM sodium chloride (tetragonal form). These crystals were of suitable size for X-ray analysis but diffracted only to 2.7 Å and 3.8 Å for the tetragonal and hexagonal crystal form respectively (Table 1).

[0055] FcγRII was expressed in *E. coli* which, besides the comparatively low production costs and the availability, has several advantages especially when the glycosylation performed by mammalian cells is not necessary for the function of the protein as in the case of FcγRII where IgG binding occurs independently of carbohydrate attachment (Sondermann et al, 1998A). In *E. coli* a homogenous product can reproducibly be generated, which is in contrast to the expression in mammalian cells where batch dependent variances are often observed. In such a system the product is for several days exposed to proteases at temperatures of more than 30°C. In contrary, the expression of the protein in *E. coli* under the control of the strong T7 promoter at 37°C frequently leads to the formation of protease inaccessible inclusion bodies. A further advantage of the expression in bacteria is that the material could be considered to be free of pathogenic germs, which might derive from employed fetal calf serum or the cell line itself. In mammalian expression particular care must be taken during the purification of the target protein because potential effective hormones or growth factors might be copurified. One case where the effects of sFcγR were ascribed to a TGFβ1 contamination is already reported (Galon et al, 1995).

1.6 Purification

[0056] The purification procedure is straightforward. It consists of three steps which can easily be performed in a single day. The protein is obtained in a pure form and in high yields and could even be obtained in considerable quality without the expensive IgG affinity column. The success of such a protocol would depend on the careful preparation of the inclusion bodies, as most of the impurities can be eliminated already in the first purification step.

1.7 Characterization

[0057] The purified FcγRIIb was characterized by SDS-PAGE and isoelectric focussing as well as N-terminal sequencing and mass spectroscopy. Thus, the material can be considered pure and homogeneous with respect to its chemical composition, but the intriguing question whether the receptor is correctly folded remains to be discussed. All cysteins are paired, since no free thiol groups are detected with Ellman's test. The material is monomeric and eludes with the expected retention time in peaks of symmetrical shape from a size exclusion chromatography column. Furthermore, FcγRIIb binds to IgG sepharose, recombinant FcγRIIb from *E. coli* is active because it specifically binds IgG.

1.8 Crystallization

[0058] The orthorhombic crystal form of FcγRIIb diffracted X-rays to a resolution of 1.7 Å, which is a drastic improvement compared to previously reported crystals of the same molecule derived from insect cell expression (Sondermann et al, 1998A). These crystals diffracted to 2.9 Å and were of space group P3₁21. Thus, the glycosylation of the insect cell derived receptor influences the crystallization conditions. Instead of the trigonal space group, three different crystal forms are found. After a possible solution of the structure these crystal forms will help identify artificial conformations of the protein due to crystal contacts.

[0059] FcγRs do not exhibit any sequence similarity to other proteins but due to a conserved cystein spacing they are affiliated to the immunoglobulin super family. Consequently, we tried to solve its structure by molecular replacement, but extensive trials using IgG domains from a variety of molecules failed. Thus the structure of FcγRIIb has to be solved by the methods of multiple isomorphous replacement.

[0060] We have shown for the first time that FcγRIIb can be obtained in an active form from *E. coli*. This is the basis for crystallographic investigations that will soon, due to the already gained crystals of exceptional quality, result in the structure solution of this important molecule. The structure will provide information on the IgG binding site and provide a starting point for the knowledge based design of drugs that interfere with recognition of the ligand by its receptor. Furthermore, because of the high homology between FcγRIIb and other FcRs including FcεRIa it seems possible that these molecules can be produced in the same way, which would provide valuable material for the ongoing research.

Example 2**2.1 Methods****5 Protein chemistry**

[0061] Recombinant soluble human FcγRIIb was expressed in E.coli, refolded purified and crystallized as described elsewhere (Sondermann et al, 1998B). Briefly, the putative extracellular region of hFcγRIIb2 (Engelhardt et al, 1990) was overexpressed in E. coli. Inclusion bodies were purified by lysozyme treatment of the cells and subsequent sonification. The resulting suspension was centrifuged (30 min 30,000 x g) and washed with buffer containing 0.5% LDAO. A centrifugation step and resuspension in LDAO containing buffer was repeated once before this procedure was repeated twice without LDAO. The inclusion bodies were solved in 6 M guanidine hydrochloride and the protein was renatured as described. The dialyzed and filtrated protein solution was applied to a hlgG sepharose column and eluted by pH jump. The concentrated neutralized fractions were subjected to size-exclusion chromatography on a Superdex-75 column (26/60, Pharmacia).

Crystallization

[0062] Crystallization was performed in sitting drops at 20°C using the vapor diffusion technique. Crystallization screens were performed by changing pH, salt, precipitant and additives. The final crystals used for data collection were grown in 33% PEG2000, 0.2 M sodium acetate, pH 5.4 (orthorhombic form) 26% PEG8000, 0.2 M sodium acetate, pH 5.6, 10% (v/v) 1,4-dioxane, 100 mM sodium chloride (tetragonal form), and 26% PEG8000, 0.2 M sodium acetate, pH 5.6, 5mM ZN(OAc)₂, 100mM sodium chloride (hexagonal form). The insect cell derived protein was crystallized in 32% PEG6000, 0.2 M sodium acetate, pH 5.3.

25 Preparation of heavy-atom derivatives

[0063] The heavy-atom derivatives were prepared by soaking the crystals in the crystallization buffer containing 2 mM platinum(II)-(2,2'-6,2''terpyridinium) chloride for 24 hours or 10 mM uranylchloride for 8 days.

30 X-ray data collection

[0064] Diffraction data was collected on an image plate system (MAR research) using graphite monochromated CuK_α radiation from a RU200b rotating anode generator (Rigaku) operated at 50 kV and 100 mA. The reflections were integrated with the program MOSFLM 5.50 (Leslie, 1997) and subsequently the data was scaled and truncated to obtain the structure-factor amplitudes using routines from the CCP4 program suite (Collaborative Computational Project, 1994).

Structure determination

[0065] The structure was solved with the standard procedures of the MIR method. From the large number of soaks carried out with different heavy-atom components only the two compounds yielded interpretable Patterson maps. The heavy-atom positions for each derivative were determined from difference Patterson maps and initial phases were calculated. Cross-phased difference Fourier maps were used to confirm heavy atom positions and establish a common origin for the derivatives. Anomalous data were included to discriminate between the enantiomers. The heavy atom parameters were further refined with the program MLPHARE from the CCP4 package leading to the statistics compiled in Table 2. An electron-density map was calculated to a resolution of 2.1 Å and the phases were improved further by solvent flattening and histogram matching with the program DM from the CCP4 suite. The resulting electron density map was of sufficient quality to build most of the amino acid residues. Model building was performed with O (Jones et al, 1991) on an Indigo2 work station (Silicon Graphics Incorporation). The structure refinement was done with XPLOR (Brünger et al, 1987) by gradually increasing the resolution to 1.7 Å using the parameter set of Engh and Huber (Engh & Huber, 1991). When the structure was complete after several rounds of model building and individual restraint B-factors refinement ($R_{\text{fac}} = 29\%$ / $R_{\text{Free}} = 36\%$), 150 water molecules were built into the electron density when a Fo-Fc map contoured at 3.5 σ coincided with well defined electron density of a 2Fo-Fc map contoured at 1 σ . The resulting refinement statistic is shown in Table 3.

2.2 Structure determination

[0066] The crystal structure of recombinant soluble human FcγRIIb was solved by multiple isomorphous replacement (MIR) to 1.7 Å resolution, since a structure solution by molecular replacement with isolated domains of the Fc fragment from human IgG1 (Huber et al, 1976, PDB entry 1fc1; Deisenhofer, 1981) failed. The putative extracellular part of the receptor (amino acid residues 1-187 as depicted in SEQ ID NO:2) was used for crystallization trials (Sondermann et al, 1998B) while the model contains the residues 5-176 as the termini are flexible and not traceable into the electron density. Additionally, the model contains 150 water molecules and the refinement statistics are summarized in Table 2. The structure contains a cis proline at position 11. None of the main chain torsion angles is located in disallowed regions of the Ramachandran plot. The fully refined model was used to solve the structure of the same protein in crystals of space group P4₂-2₁ and of the glycosylated form derived from insect cells in crystals of space group P3₁21 (Table 2).

[0067] The polypeptide chain of FcγRIIb folds into two Ig-like domains as expected from its affiliation with the immunoglobulin super family. Each domain consists of two beta sheets that are arranged in a sandwich with the conserved disulfide bridge connecting strands B and F on the opposing sheets (Fig. 3). Three anti-parallel β-strands (A1, B, E) oppose a sheet of 5 β-strands (C', C, F, G, A2), whereby strand A1 leaves the 3-stranded β-sheet and crosses over to the 4-stranded anti-parallel sheet adding the short parallel 5th strand A2. The arrangement of secondary structure elements as well as their connectivity is identical in both domains of the FcγRIIb and a rigid body fit of one domain onto the other revealed a r.m.s. distance of 1.29 Å of 67 matching Cα atoms.

[0068] The domains are arranged nearly perpendicularly to each other enclosing an angle of 70 degrees between their long axes forming a heart-shaped overall structure. This arrangement results in an extensive contact region between the domains (Fig. 4). Residues from strand A2 and from the segment linking A2 and A1 of the N-terminal domain intermesh with residues of strands A1 and B from the C-terminal domain. This region is tightly packed and the interaction is strengthened by several hydrogen bonds resulting in a rigid arrangement. This is confirmed by the conservation of the structure in three different space groups. In orthorhombic, tetragonal and hexagonal (insect cell derived) crystal forms a deviation of less than 2° in the interdomain angle is found.

2.3 Overall structures

[0069] The structure of recombinant human FcγRIIb derived from E.coli was solved by MIR to 1.7 Å resolution from orthorhombic crystals. An essentially identical structure is found in tetragonal and with protein derived from insect cells in hexagonal crystals. In all three structures the last nine residues of the polypeptide chain were found disordered. The flexibility of the C-terminal linker region between the structured core of the molecule and the transmembrane part may be functionally relevant to allow some reorientation of the receptor to enhance the recognition of the Fc parts in immunocomplexes.

2.4 Homologue receptors

[0070] The Ig domains found in the Ig super family of proteins are characterized by a beta sandwich structure with a conserved disulfide bridge connecting two strands of the opposing sheets. The typical arrangement of 3 and 4 anti parallel beta strands that form a sandwich as found in FcγRIIb occurs also in the T cell receptor, Fc fragment, CD4 or the Fab fragment. A structural alignment of the individual Ig domains of these molecules with the two domains of FcγRIIb shows a common, closely related structure. The relative arrangement of the domains, however, is not related in these molecules and covers a broad sector. Despite the structural similarity between Ig domains from different molecules and the strikingly low r.m.s. deviation of Cα atoms that result when the two domains of FcγRII are superimposed, no significant sequence similarity is found (Figs. 5a and 5b). A structure-based sequence alignment shows a conserved hydrophobicity pattern along the sequence of the domains, together with, beside the cysteins, only few identical amino acid residues. We first prepared a structure-based alignment of the two C-terminal domains of the IgG1 heavy chain and the FcγRIIb and added the sequences of the other related FcγRI and the FcεRIa domains. This shows that the sequences of the three domain FcγRI and the two domain receptors are compatible with the hydrophobicity pattern of Ig domains and several conserved amino acid residues are revealed. Firstly, the different domains of an FcR are more related to each other than to Ig domains from other molecules of the Ig super family. Secondly, the N-terminal domains of the receptors relate to each other as the second domains do. Thirdly, the sequence of the third domain of FcγRI shows features from both groups of domains. Taken together, we confirm the affiliation of the FcRs to the Ig super family and speculate that all FcR-domains originate from a common ancestor, an ancient one domain receptor that acquired a second domain by gene duplication. Further divergent development of such a two domain receptor resulted in the present diversity, including FcγRI that acquired a third domain.

[0071] Conservation of these amino acid residues that contribute to the interdomain contact in FcγRIIb in the align-

ment are a hint to a similar domain arrangement in different receptors. In Table 4 the residues contributing with their side chains to the interdomain contact (Fig. 4) are compiled for Fc γ R11b together with the corresponding amino acid residues in other receptors according to the structure-based sequence alignment of Fig. 5b. Except for Asn15, which is not conserved between the FcRs, the involved residues are identical or conservatively replaced providing strong support for a similar structure and domain arrangement in all FcRs.

2.5 The contact interface to IgG

[0072] Limited information about the interactions of FcRs with their ligands is available from mutagenesis studies (Hogarth et al, 1992; Hulett et al, 1994; Hulett et al, 1995). By systematically exchanging loops between the β -strands of Fc γ R11a for Fc ϵ R1a amino acid residues the B/C, C/E and F/G loops of the C-terminal domain were evaluated as important for ligand binding (Fig. 3, Fig. 5b). In the structure model these loops are adjacent and freely accessible to the potential ligand. Additionally, most of the amino acid residues in these loops were exchanged for alanines by single site mutations which resulted in a drastic alteration of the affinity of Fc γ R11a to dimeric human IgG1. Also, the single amino acid exchange Arg 131 to His in the C-terminal domain (C/E loop) in the high responder/low responder polymorphism, which alters the affinity of the Fc γ R11a to murine IgG1, points to that region. Thus, the amino acid residues in this area are either important for ligand binding or the structural integrity of that region. Here, the structure shows a clustering of the hydrophobic amino acid residues Pro 114, Leu 115 and Val 116 in the neighbourhood of Tyr 157. This patch is separated from the region Leu 159, Phe 121 and Phe 129 by the positively charged amino acid residues Arg 131 and Lys 117 which protrude from the core structure (Fig. 5b).

2.6 Glycosylation

[0073] In the sequence of Fc γ R11b three potential N-glycosylation sites are found. All three sites are on the surface of the molecule and are accessible. They are located in the E/F loops (N61 and N142) of both domains and on strand E (N135) of the C-terminal domain (Fig. 3, Fig. 6). Since the material used for the solution of this structure was obtained from *E. coli*, it does not contain carbohydrates, while the FcRs isolated from mammalian cells are highly glycosylated. The three potential glycosylation sites are located rather far from the putative IgG binding region, and non-glycosylated Fc γ R11b binds human IgG, suggesting a minor role of glycosylation in binding. This was confirmed by the structure of the Fc γ R11b produced in insect cells which is glycosylated (Sondermann et al, 1998A). Except for a 2° change of the interdomain angle possibly due to different crystal contacts, no differences between the glycosylated and unglycosylated protein structures were found. The three glycosylation sites are only optionally used as shown by SDS-PAGE where the material appears in 4 bands. No additional electron density for those sugars was found a consequence of chemical and structural heterogeneity.

2.7 The modeled complex

[0074] The newly solved structure of Fc γ R11b complements the information gained from the structure of the Fc fragment and the available biochemical data regarding the Fc γ R:IgG complex.

[0075] While diverse biochemical information concerning the binding site of Fc γ R11b (see above) is available, only limited data exists regarding the contact area contributed by the antibody. The IgG isotypes are closely related and exhibit graded affinities to Fc γ Rs. However, they still carry too many amino acid exchanges for the determination of the binding site and the preparation of IgG mutants is tedious. The only available information results from experiments with Fc γ R bearing cells on which bound immunocomplexes could be displaced with protein A (Ades et al, 1976), suggesting an at least partially overlapping binding site of protein A and Fc γ R11b on the antibody.

[0076] With the structures of both constituents at hands we attempted to model the Fc γ R11:IgG complex using the program FTDock (Gabb et al, 1997). FTDock uses Fourier correlation theory for evaluation of the shape and electrostatic complementarity of the complex component surfaces. In the hands of the authors the program has produced good results in predicting complex structures, but in some cases additional biochemical information on the location of the contact area was needed to exclude false positive solutions.

[0077] Without applying additional restrictions concerning the region of the contact surface between Fc γ R11b and the Fc fragment, the calculations resulted in a single solution clearly scoring above the rather constant background. The program predicted a complex structure with the B/C, C/E and F/G loops of the Fc γ R11b domain 2 contributing to the contact site as predicted by the mutagenesis experiments. The only observed interaction of the N-terminal domain with the Fc fragment is via E19 that forms a salt bridge to a lysine of the CH₂ domain. Some involvement of residues of the N-terminal domain in complex formation is expected since the N-terminal domain of Fc γ R11a cannot be exchanged against the corresponding domain of Fc ϵ R1a (Hulett et al, 1995) without losing the ligand binding capability of the receptor.

[0078] From the predicted interaction a model of the membrane bound complex between IgG and Fc γ R11b is pro-

posed (Fig. 7). Two FcγRIIb bind into the cleft between the third and the fourth domain of the IgG heavy chains employing the 2-fold symmetry of the Fc fragment. Protein A (Deisenhofer et al, 1978; Deisenhofer, 1981) as well as protein G (Sauer-Eriksson et al, 1995) and the neonatal FcR (Burmeister et al, 1994) bind to a surface region around the exposed hydrophobic residue Ile 253 of the Fc fragment. FcγRIIb binds to a region in the vicinity consistent with the competitive binding of protein A and FcγR to the antibody. The 2:1 stoichiometry between FcγRIIb and Fc fragment in the complex could be shown in equilibrium gel filtration experiments (Sondermann et al, 1998A).

[0079] The complex can be positioned upright on the membrane, with the truncated C-termini of FcγRIIb oriented towards the membrane. The N-terminal domain of the receptor lies parallel to the membrane between the Fab arms when the complex is viewed along the Fc fragment. If the FcRs have evolved from a common one domain receptor we expect that the amino acid residues of the N-terminal domain that correspond to the binding region of the C-terminal domain form a second putative binding site. The corresponding surface region is accessible in the proposed complex and forms a large uncharged patch with a hydrophobic ridge comprising amino acid residues Pro 47, Leu 45, Phe 40, Leu 75, Pro 3, Pro 2 and Ala 1 (Fig. 6b). This region might represent a binding site for other ligands that have been discussed for FcγRIIb to explain the signalling capabilities of its soluble form.

[0080] Thus the modeled complex structure is consistent with the available biochemical data.

Fig. 1: 15% reducing SDS PAGE showing the purification of sFcγRIIb

Lane 1: Molecular weight marker. Lane 2: E. coli lysate before induction. Lane 3: E. coli lysate 1 h after induction. Lane 4: E. coli lysate 4 h after induction. Lane 5: Purified inclusion bodies of sFcγRIIb. Lane 6: Eluate of the hlgG affinity column. Lane 7: Pooled fractions of the gel filtration column.

Fig. 2: Equilibrium gel filtration

1 μg hFc solved in 10 μl equilibration buffer (10 μg sFcγRIIb/ml PBS) was applied to a size exclusion chromatography column and the absorbance of the effluent was measured (280 nm) as a function of time. The injected Fc fragment forms a complex with the sFcγRIIb in the equilibration buffer (t = 22 min). The negative peak of consumed sFcγRIIb is observed at t = 26 min.

Fig. 3: Overall structure of human sFcγRIIb

Stereo ribbon representation of the sFcγRIIb structure. The loops supposed to be important for IgG binding are depicted in red with some of the residues within the binding site and the conserved disulfide bridge in ball and stick representation. The potential N-glycosylation sites are shown as green balls. The termini are labeled and the β-strands are numbered consecutively for the N-terminal domain in black and for the C-terminal domain in blue. The figure was created using the programs MOLSCRIPT (Kraulis, 1991) and RENDER (Merritt and Murphy, 1994).

Fig. 4: Interdomain contacts

The figure shows a close-up on the residues involved in the interdomain contacts of sFcγRIIb. The amino acid residues of the N-terminal domain are depicted blue and the residues of the C-terminal domain yellow. The model is covered by a 2Fo-Fc electron density contoured at 1 σ obtained from the final coordinates. Hydrogen bridges between the domains are represented by white lines. The figure was created using the program MAIN (Turk, 1992).

Fig. 5a: Superposition of the two FcγRIIb domains and the CH2 domain of human IgG1

Both domains of FcγRIIb and the CH2 domain of hlgG1 were superimposed. The N-terminal domain is depicted in blue, the C-terminal domain in red and the CH2 domain of hlgG1 in green. The respective termini are labeled and the conserved disulfide bridges are depicted as thin lines.

Fig. 5b: Structure based sequence alignment of the sFcγRIIb domains with domains of other members of the FcR family

The upper part of the figure shows the structure based sequence alignment of the FcγRIIb and hlgG1 Fc fragment domains performed with the program GBF-3D-FIT (Lessel & Schomburg, 1994). Amino acid residues with a Cα distance of less than 2.0 Å in the superimposed domains are masked: lilac for matching residues between the Fc fragment domains; yellow for residues in the FcγRIIb domains; and green when they can be superimposed in all four domains. The β-strands are indicated below this part of the alignment and are labeled consistent with Figure 3. The lower part of the figure shows the alignment of the amino acid sequences from the other FcγRs and the homologue FcεRIa to the profile given in the upper part of the figure using routines from the GCG package (Genetics Computer Group, 1994). The upper and lower row of numbering refer to the N- and C-terminal domains of FcγRIIb. The conserved cysteines are typed in magenta and the potential glycosylation sites in blue. Identical residues within the first domain are masked orange, those in the second domain pink and green when the residues are conserved within both domains. The less conserved third domain of FcγRI is aligned between the first and the second

domains. Red arrows point to residues that are involved in side chain contacts between the first and the second domain while blue arrows depict residues that are relevant for IgG binding. The figure was produced with the program ALSCRIPT (Barton, 1993).

Fig. 6: The putative binding sites of FcγRIIb

Solid surface representations of FcγRIIb as produced with GRASP (Nicholls et al, 1991), the color coding is according to the relative surface potential from negative (red) to positive (blue). Fig. 6a shows the molecule as in Fig. 3 by a rotation of about 90° counter-clockwise around the vertical. In Fig. 6b the molecule is rotated 90° clockwise around the same axis. Both views show the putative binding regions on the C-terminal (Fig. 6a) and the N-terminal domain (Fig. 6b). The amino acid residues discussed in the text are labeled.

Fig. 7: Model of the FcγR-IgG complex

The cartoon shows a complete complex of two FcγRIIb binding one antibody as suggested by the program FFT-DOCK. The heavy chains of the antibody are depicted in red and green and the light chains in yellow. The blue atoms represent the C-terminal domain of sFcγRIIb while the white ones represent the N-terminal domain. A blue column connects the receptor to the membrane instead of the flexible linker region that remained invisible in the electron density. The image was produced with the program POVray.

Fig. 8: Alignment of the amino acid sequence of the extracellular parts of FcγR and FcεRIa

Figure 8 shows an alignment of amino acid sequences of the extracellular parts of some FcγRs and FcεRI.

Table 1

Crystallographic results			
The obtained preliminary crystallographic data are shown in this table.			
	Orthorhombic	Tetragonal	Hexagonal
Space group	P2 ₁ 2 ₁ 2 ₁ [19]	P4 ₂ 2 ₁ 2 [94]	P3 [143]
Unit cell dimensions	a=40.8Å, b=50.9Å, c=80.5Å, α=90°, β=90°, γ=90°	a=85.7Å, b=85.7Å, c=63.4Å, α=90°, β=90°, γ=90°	a=80.9Å, b=80.9Å c=157.0Å, α=90°, β=90°, γ=90°
R _{merge}	5.8%	9.8%	13.6%
Resolution	1.7Å	2.7Å	3.8Å
Unique	18,040	6,616	7,210
Completeness	89.1%	97.1%	63.0%
Multiplicity	3.5	4.4	1.3
V _M , molecules per asymmetric unit, solvent content	2.09Å ³ /Da, 1 mol., 41 % solvent	2.91Å ³ /Da, 1 mol, 58% solvent	2.97Å ³ /Da, 5 mol, 59% solvent

Table 2: Data collection statistics

Derivative	Space Group	No. of unique reflections	Multiplicity	Resolution (Å)	Completeness (overall/last shell) (%)	R _m (%)	No. of sites	Phasing power
NATI	P2 ₁ 2 ₁ 2 ₁	18009	3.6	1.74	92.9/86.4	6.5		
NATI	P4 ₂ 2 ₁ 2	6615	4.5	2.70	97.1/94.3	10.1		
NATI-Baculo	P3 ₁ 21	3545	2.5	3.0	93.0/98.9	14.4		
UOAc	P2 ₁ 2 ₁ 2 ₁	7722	4.2	2.1	96.8/95.7	7.3	1	1.79
PtPy	P2 ₁ 2 ₁ 2 ₁	5520	3.9	2.3	89.7/49.6	10.5	1	1.39

$$R_m = \Sigma |I_h - \langle I_h \rangle| / \Sigma \langle I_h \rangle$$

Phasing power: $\langle F_H \rangle / E$, where $\langle F_H \rangle = \Sigma (F_H^2 / n)^{1/2}$ is the r.m.s. heavy atom structure amplitude.

$E = \Sigma [(F_{PHC} - F_{PH})^2 / n]^{1/2}$ is the residual lack of closure error with F_{PH} being the structure factor amplitude and $F_{PHC} = |F_P + F_H|$ the calculated structure factor amplitude of the derivative.

Table 3

Refinement statistics	
Resolution range (Å)	8.0 - 1.74 Å
No. of unique reflections ($F > O\sigma(F)$)	16252
R factor	19.4
R_{free}^*	27.9
No. of atoms per asymmetric unit	
protein	1371
solvent	150
Rms deviation from ideal geometry	
bond length (Å)	0.009
bond angle (°)	2.007
Average B factors (Å ²)	
protein main chain	18.8
protein side chain	25.2
solvent	36.7
Rms deviation of bonded B factors (Å ²)	4.1

* R_{free} : 5% of the reflections were used as a reference data set and were not included in the refinement.

Table 4

Residues that contribute to the interdomain contact via side chains				
FcγRIIb	FcγRIIa	FcγRIII	FcγRI	FcεRIa
Asn15	Asn	Ser	Ser	Arg
Asp20	Asp	Asp	Glu	Glu
Gln91	Gln	Gln	Gln	Gln
His108	His	His	His	His
Trp110	Trp	Trp	Trp	Trp

SEQ ID NO:1

5

GTPAAPPKAV LRLEPQWINV LQEDSVTLTC RGTQSPESDS IQWFHNGNLI PTHTQPSYRF
 KANNNDSGEY TCQTGQTSVS DPVHLTVLSE WVLQTPHLE FQGETIVLR CHSWKDKPLV
 KVTFFQNGKS KKFSRSDPNF SIPQANHSYS GDYHCTGNIG YTLYSSKPVT ITVQAP...S
 SSPMGII

10

SEQ ID NO:2

15

AVPQKPK VSLNPPWNRI PKGENVTLC NGNNFFEVS TKWFHNGSL EETNSSLNIV
 NAKFEDSGEY KCQHQQVNES EPVYLEVFS WLLQASAEV VMEGQPLFLR CHGWRNWDVY
 KVIYYKDGEA LKYWYENHNI SITNATVEDS GTYYCTGKVV QLDYESEPLN ITVIKAPREK
 YWLQ(F)

20

25

SEQ ID NO:3

30

aacagaa ttccATGggg acacotgcag
 181 ctcccccaaa ggctgtgctg aaactcgagc cccagtggat caactgtgctc caggaggact
 241 ctgtgactct gacatgccgg gggactcaca gccctgagag cgaactcatt cagtgggtcc
 301 acaatgggaa tctcattccc acccacagc agcccagcta caggttcaag gccacaaca
 361 atgacagcgg ggagtacacg tgccagactg gccagaccag cctcagcgac cctgtgcatc
 421 tgacagtgtt ttctgagtgg ctggtgtctc agaccctca cctggagttc caggagggag
 481 aaaccatcgt gctgaggtgc cacagctgga aggacaagcc tctggtcaag gtcacattct
 541 tccagaatgg aaaatccaag aaattttccc gtteggatcc caacttctcc atcccacaag
 601 caaaccacag tcacagtggg gattaccatt gcacaggaaa cataggctac acgctgtact
 661 catccaagcc tgtgaccatc actgtccaag ctcccagctc ttcaccgatg gggatcattT
 721 AGgctgtcga gactggg

40

45

50

55

SEQ ID NO:4

5
121 cagaaaccta aggtctcctt gaacctcca tggaatagaa tatttaaagg agagaatgtg
181 actcttacat gtaatgggaa caatttcttt gaagtcagtt ccaccaaatg gtccacaat
241 ggcagccttt cagaagagac aaattcaagt ttgaatattg tgaatgccaa atttgaagac
10 301 agtggagaaat acaaatgtca gcaccaataa gttaatgaga gtgaacctgt gtacctggaa
361 gtcttcagtg actggctgct ccttcaggcc tctgctgagg tggatgagg gggccagccc
421 ctcttcctca ggtgccatgg ttggaggaaac tggatgtgt acaaggtgat ctattataag
481 gatggtgaag ctctcaagta ctggtatgag aaccacaaca tctccattac aaatgccaca
15 541 gttgaagaca gtggaacctc ctactgtacg ggcaaatgtt ggcagctgga ctatgagctc
601 gagccctcca acattactgt aataaaagct cgcgtgaga agtactggct acaattttag
661 gattccattg

20

SEQ ID NO:5

human FcγRIIb2
26 1 ggctgtgact gctgtgctct gggcgccact cgctccaggg agtgatggga atcctgtcat
61 ttttacctgt ccttgccact gagagtact gggctgactg caagtcccc cagccttggg
121 gtcatatgct tctgtggaca gctgtgctat tcttggtctc tgttgctggg acacctgcag
181 ctccccaaa ggcgtgtgctg aaactcgagc ccagtggtat caactgtctc caggaggact
241 ctgtgactct gacatgccgg gggactcaca gccctgagag cgactccatt cagtggttcc
30 301 acaatgggaa tctcattccc acccacagcg agcccagcta caggttcaag gccacaaca
361 atgacagcgg ggagtacacg tgccagactg gccagaccag cctcagcgac cctgtgcatc
421 tgacagtgtc ttctgagtgg ctggtgctcc agacccctca cctggagtcc caggagggag
481 aaaccatcgt gctgaggtgc cacagctgga aggacaagcc tctggtcaag gtcacattct
35 541 tccagaatgg aaaatccaag aaattttccc gtteggatcc caacttctcc atcccacaag
601 caaaccacag tcacagtggg gattaccatt gcacaggaaa cataggctac acgctgtact
661 catccaagcc tgtgaccatc actgtccaag ctcccagctc ttcaccgatg gggatcattg
721 tggctgtggt cactgggatt gctgtagctg ccattgttgc tgcgttagtg gccttgatct
40 781 actgcaggaa aaagcggatt tcagccaatc cactaatcc tgatgaggct gacaaagtgc
841 gggctgagaa cacaatcacc tattcacttc tcatgcaccc ggatgctctg gaagagcctg
901 atgaccagaa ccgtattttg tctccattgt ctgcatggg gatttgagaa gaaatcagag
961 agggaagatc tggattttcc tggcctaaat tccccttggg gaggacaggg agatgctgca
45 1021 gttccaaaag agaaggttcc ttccagagtc atctacctga gtcttgaagc tccctgtcct
1081 gaaagccaca gacaatatgg tcccgaatgc ccgactgcac cttctgtgct tcagctcttc
1141 ttgacatcaa ggctcttccg ttccacatcc acacagccaa tccaattaat caaaccactg
1201 ttattaacag ataatagcaa cttgggaaat gcttatgtta cagggttacgt gagaacaatc
50 1261 atgtaaatct atatgatttc agaaatgtta aaatagacta acctctacca gcacattaaa
1321 agtgattgtt tctgggtgat aaaattattg atgattttta tttctttat tttctataa
1381 agatcatata ttacttttat aataaaacat tataaaaac

55

SEQ ID NO:6

5

human FcεRIα

1 agatctcagc acagtaagca ccaggagtcc atgaagaaga tggctcctgc catggaatcc
61 cctactctac tgtgtgtagc cttactgttc ttcgctccag atggcgtggt agcagtcctt
10 121 cagaaacctc aggtctcctt gaacctcca tggaatagaa tatttaaagg agagaatgtg
181 actcttacat gtaatgggaa caatttcttt gaagtcagtt ccaccaaatg gttccacaat
241 ggcagccttt cagaagagac aaattcaagt ttgaatattg tgaatgccaa atttgaagac
301 agtggagaat acaaatgtca gcaccaacaa gttaatgaga gtgaacctgt gtacctggaa
15 361 gtcttcagtg actggctgct ccttcaggcc tctgctgagg tggatgagga gggccagccc
421 ctcttcctca ggtgccatgg ttggaggaac tgggatgtgt acaagggtgat ctattataag
481 gatgggtgaag ctctcaagta ctggtatgag aaccacaaca tctccattac aaatgccaca
541 gttgaagaca gtggaacctc ctactgtacg ggcaaagtgt ggcagctgga ctatgagtct
20 601 gageccctca acattactgt aataaaagct ccgctgaga agtactggct acaatttttt
661 atcccattgt tgggtggtgat tctgtttgct gtggacacag gattatttat ctcaactcag
721 cagcaggtca catttctctt gaagattaag agaaccagga aaggcttcag acttctgaac
781 ccacatccta agccaaaccc caaaaacaac tgatataatt aactcaagaa atatttgcaa
25 841 cattagtttt ttccagcat cagcaattgc tactcaattg tcaaacacag cttgcaatat
901 acatagaaac gtctgtgctc aaggatttat agaaatgctt cattaaactg agtgaaactg
961 attaagtggc atgtaatagt aagtgtcaa ttaacattgg ttgaataaat gagagaatga
1021 atagattcat ttattagcat ttgtaaaaga gatgttcaat ttagatct

30

35

40

45

50

55

SEQ ID NO:7

5

human mRNA for high affinity Fc receptor (FcγRI)

10

15

20

25

30

35

40

45

50

55

1 gacagatttc actgctccca ccagcttggg gacaacatgt ggttcttgac aactctgctc
61 ctttgggttc cagttgatgg gcaagtggac accacaaagg cagtgatctc ttgcagcct
121 ccatgggtca gcgtgttcca agaggaaacc gtaaccttgc actgtgaggt gctccatctg
181 cctgggagca gctctacaca gtggtttctc aatggcacag ccactcagac ctgcaccccc
241 agctacagaa tcacctctgc cagtgtcaat gacagtgggt aatacagggt ccagagaggt
301 ctctcagggc gaagtgacct catcacagctg gaaatccaca gaggctgggt actactgcag
361 gtctccagca gagtcttcac ggaaggagaa cctctggcct tgagggtgca tgcgtggaag
421 gataagctgg tgtacaatgt gctttactat cgaaatggca aagcctttaa gtttttccac
481 tgggaattcta acctcaccat tctgaaaacc aacataagtc acaatggcac ctaccattgc
541 tcaggcatgg gaaagcatcg ctacacatca gcaggaatat ctgtcactgt gaaagagcta
601 tttccagctc cagtgtgaa tgcattctgt acatccccac tctggagggt gaactctggtc
661 accctgagct gtgaaacaaa gttgctcttg cagaggcctg gtttgagct ttacttctcc
721 ttctacatgg gcagcaagac cctgcgaggt aggaacacat cctctgaata ccaaatacta
781 actgctagaa gagaagactc tgggttatac tgggtgcgag ctgccacaga ggatggaaat
841 gtctttaagc gcagccctga gttggagctt caagtgtctg gcctccagtt accaactcct
901 gtctgggttc atgtcctttt ctatctggca gtgggaataa tgtttttagt gaacactggt
961 ctctgggtga caatacgtaa agaactgaaa agaaagaaaa agtgggattt agaaatctct
1021 ttggattctg gtcattgaga gaaggtaact tccagccttc aagaagacag acatttagaa
1081 gaagagctga aatgtcagga acaaaaagaa gaacagctgc aggaaggggt gcaccggaag
1141 gagccccagg gggccacgta gcagcggctc agtgggtggc catcgatctg gaccgtcccc
1201 tgcccacttg cccccgtga gcaactgcgt caaacatcca aaagttcaac aacaccagaa
1261 ctgtgtgtct catggtatgt aactcttaa gcaaataaat gaactgactt caaaaaaaaa
1321 a

SEQ ID NO:8

5

human FCγRIIa

10

15

20

25

30

35

40

45

50

55

1 cccaaatgtc tcagaatgta tgtcccagaa acctgtggct gcttcaacca ttgacagttt
 61 tgctgctgct ggcttctgca gacagtcaag ctgcagctcc cccaaaggct gtgctgaaac
 121 ttgagccccc gtggatcaac gtgctccagg aggactctgt gactctgaca tgccaggggg
 181 ctcgcagccc tgagagcgac tccattcagt ggttccacaa tgggaatctc attcccaccc
 241 acacgcagcc cagctacagg ttcaaggcca acaacaatga cagcggggag tacacgtgdc
 301 agactggcca gaccagcctc agcgaccctg tgcactctgac tgtgctttcc gaatggctgg
 361 tgctccagac ccttcacctg gagttccagg agggagaaac catcatgctg aggtgccaca
 421 gctggaagga caagcctctg gtcaaggcca cattcttcca gaatggaaaa tcccagaaat
 481 tctcccgttt ggatcccacc ttctccatcc cacaagcaaa ccacagtcac agtgggtgatt
 541 accactgcac aggaacata ggctacacgc tgttctcacc caagcctgtg accatcactg
 601 tccaagtgcc cagcatgggc agctcttcac caatggggat cattgtggct gtggtcattg
 661 cagctgctgt agcagccatt gttgctgctg tagtggcctt gatctactgc agggaaaagc
 721 ggatttcagc caattccact gatcctgtga aggtgcccc atttgagcca cctggacgtc
 781 aaatgattgc catcagaaag agacaacttg aagaaaccaa caatgactat gaaacagctg
 841 acggcggcta catgactctg aaccccaggg cacctactga cgatgataaa aacatctacc
 901 tgactcttcc tcccacgcac catgtcaaca gtaataacta aagagtaacy ttatgccatg
 961 tggtcatact ctgagcttgc tgatggatga caaaaagagg ggaattgtta aaggaaaatt
 1021 taaatggaga ctgaaaaaat cctgagcaaa caaaaccacc tggcccttag aaatagcttt
 1081 aactttgctt aaactacaaa cacaagcaaa acttcacggg gtcatactac atacaagcat
 1141 aagcaaaact taacttggat catttctggt aaatgcttat gttagaata agacaacccc
 1201 agccaatcac aagcagccta ctaacatata attaggtgac tagggacttt ctaagaagat
 1261 acctaccccc aaaaaacaat tatgtaattg aaaaccaacc gattgccttt attttgettc
 1321 cacattttcc caataaatac ttgcctgtga cattttgccg ctggaacact aaacttcag
 1381 aattgcgect cagatttttc ctttaacatc tttttttttt ttgacagagt ctcaatctgt
 1441 taccagggtt ggagtgcagt ggtgctatct tggtcactg caaacccgcc tcccagggtt
 1501 aagcgattct tatgcctcag cctcccagta gctgggatta gaggcagtgt ccatcatacc
 1561 cagctaattt ttgtattttt ttttttttat ttttagtaga gacagggttt cgcaatgttg
 1621 gccaggccga tctcgaactt ctggcctcta gcgatctgcc cgcctcggcc tccc aaagt
 1681 ctgggatgac cgcctcagcc ccaatgtcca gcctctttaa catcttcttt cctatgcctt
 1741 ctctgtggat cctactgct ggtttctgcc ttctccatgc tgagaacaaa atcacctatt
 1801 cactgcttat gcagtcggaa gctccagaag aacaaagagc ccaattacca gaaccacatt
 1861 aagtctccat tgttttgcct tgggatttga gaagagaatt agagaggtga ggatctggta
 1921 tttctgggac taaattccct tggggaagac gaagggatgc tgcagttcca aaagagaagg
 1981 actcttccag agtcatctac ctgagtcacca aagctccctg tctgaaaagc cacagacaat
 2041 atgggtccaa atgactgact gcacctctg tgcctcagcc gttcttgaca tcaagaatct
 2101 tctgttccac atccacacag ccaatacaat tagtcaaacc actgttatta acagatgtag
 2161 caacatgaga aacgcttatg ttacagggtta catgagagca atcatgtaag tctatatgac
 2221 ttcagaaatg ttaaaataga ctaacctcta acaacaaatt aaaagtgatt gtttcaaggt
 2281 gatgcaatta ttgatgacct attttatatt tctataatga tcatatatta cttttgtaat
 2341 aaaacattat aaccaaacc

SEQ ID NO:9

5

human FcγRIII

1 tctttggtga cttgtccact ccagtgtggc atcatgtggc agctgtcct cccaactgct
 61 ctgctacttc tagtttcage tggcatgcgg actgaagatc tcccaaaggc tgtggtgttc
 10 121 ctggagcctc aatggtacag cgtgcttgag aaggacagtg tgactctgaa gtgccaggga
 181 gctactccc ctgaggacaa ttccacacag tggtttcaca atgagagcct catctcaagc
 241 caggcctcga gctacttcat tgacgctgcc acagtcaacg acagtggaga gtacagggtg
 301 cagacaaacc tctccaccct cagtgacccg gtgcagctag aagtcacat cggtcggtg
 15 361 ttgctccagg cccctcgggt ggtgttcaag gaggaagacc ctattcacct gaggtgtcac
 421 agctggaaga aactgtctct gcataaggtc acatatttac agaatggcaa agacaggaag
 481 tattttcctc ataattctga cttccacatt ccaaaagcca cactcaaaga tagcggtctc
 541 tactttctgca gggggcttgt tgggagtaaa aatgtgtctt cagagactgt gaacatcac
 20 601 atcactcaag gtttggcagt gtcaaccatc tcatcattct ctccacctgg gtaccaagtc
 661 tctttctgct tggatggtt actccttttt gcagtggaca caggactata tttctctgtg
 721 aagacaaaca tttgaagctc aacaagagac tgggaaggacc ataaacttaa atggagaaag
 781 gacctcaag acaaatgacc cccatcccat gggagttaata agagcagtgg cagcagcatc
 25 841 tctgaacatt tctctggatt tgcaaccca tcatctcag gctctctc

30 References

[0081]

- 35 Ades, E.W., Phillips, D.J., Shore, S.L., Gordon, D.S., LaVia, M.F., Black, C.M., Reimer, C.B. (1976), Analysis of mononuclear cell surfaces with fluoresceinated Staphylococcal protein A complexed with IgG antibody or heat-aggregated γ-globulin, *J. Immunol.* 117, 2119-?.
- 40 Amigorena, S., Bonnerot, C., Drake, J.R., Choquet, D., Hunziker, W., Guillet, J.G., Webster, P., Sautes, C., Mellman, I., Fridman, W.H. (1992), Cytoplasmic domain heterogeneity and functions of IgG Fc receptors in B lymphocytes, *Science* 256, 1808-1812.
- Barton, G.C. (1993), ALSCRIPT: tool to format multiple sequence alignments, *Prot. Eng.* 6, 37-40.
- 45 Bazil, V. and Strominger, J.L. (1994), Metalloprotease and serine protease are involved in cleavage of CD43, CD44, and CD16 from stimulated human granulocytes, *J. Immunol.* 152, 1314-1322.
- Brünger, A.T., Kuriyan, J., Karplus, M. (1987), Crystallographic R factor refinement by molecular dynamics, *Science* 35, 458-460.
- 50 Burmeister, W.P., Huber, A.H., Bjorkman, P.J. (1994), Crystal structure of the complex of rat neonatal Fc receptor with Fc, *Nature* 372, 379-383.
- Ceuppens, J.L., Baroja, M.L., van Vaeck, F., Anderson, C.L. (1988), Defect in the membrane expression of high affinity 72kD Fcγ receptors on phagocytic cells in four healthy subjects, *J. Clin. Invest.* 82, 571-578.
- 55 Collaborative computational project, Number 4 (1994), The CCP4 suite: Programs for protein crystallography, *Acta crystallogr. D* 50, 760-763.

- Deisenhofer, J., Jones, T.A., Huber, R., Sjodahl, J., Sjoquist, J. (1978), Crystallization, crystal structure analysis and atomic model of the complex formed by a human Fc fragment and fragment B of protein A from *Staphylococcus aureus*, *Z. Phys. Chem.* 359, 975-985.
- 5 Deisenhofer, J. (1981), Crystallographic refinement and atomic models of a human Fc fragment and its complex with fragment B of protein A from *Staphylococcus aureus* at 2.9- and 2.8Å resolution, *Biochemistry* 20, 2361-2370.
- Dulau, L., Cheyrou, A., Aigle, M. (1989), Directed mutagenesis using PCR, *Nucleic Acids Res.* 17, 2873.
- 10 Ellman (1959), Tissue sulfhydryl groups, *Arch. Biochem. Biophys.* 82, 79-77.
- Engelhardt, W., Geerds, C., Frey, J. (1990), Distribution, inducibility and biological function of the cloned and expressed human β Fc receptor II, *Eur. J. Immunol.* 20, 1367-1377.
- 15 Engh, R.A. and Huber, R. (1991), Accurate bond and angle parameters for X-ray protein structure refinement, *Acta crystallogr.* A47, 392-400.
- Fleit, H.B., Kobasiuk, C.D., Daly, C., Furie, R., Levy, P.C., Webster, R.O. (1992), A soluble form of Fc γ RIII is present in human serum and other body fluids and is elevated at sites of inflammation, *Blood* 79, 2721-2728.
- 20 Fridman, W.H., Bonnerot, C., Daeron, M., Amigorena, S., Teillaud, J.-L., Sautès, C. (1992), Structural bases of Fc γ receptor functions, *Immunol. Rev.* 125, 49-76.
- Fridman, W.H., Teillaud, J.-L., Bouchard, C., Teillaud, C., Astier, A., Tartour, E., Galon, J., Mathiot, C., Sautès, C. (1993), Soluble Fc γ receptors, *J. Leukocyte Biol.* 54, 504-512.
- 25 Gabb, H.A., Jackson, R.M., Steinberg, M.J.E. (1997), Modelling protein docking using shape complementarity, electrostatics and biochemical information, *J. Mol. Biol.* 272, 106-120.
- Galon, J., Bouchard, C., Fridman, W.H., Sautès, C. (1995), Ligands and biological activities of soluble Fc γ receptors, *Immunol. Lett.* 44, 175-181.
- 30 Genetics Computer Group (1994), Program Manual for the Wisconsin Package Version 8, Madison, Wisconsin.
- 35 Grodberg, J. and Dunn, J.J. (1988), OmpT encodes the *Escherichia coli* outer membrane protease that cleaves T7 RNA polymerase during purification, *J. Bacteriol.* 170, 1245-1253.
- Hogarth, P.M., Hulett, M.D., Ierino, F.L., Tate, B., Powell, M.S., Brinkworth, R.I. (1992), Identification of the immunoglobulin binding regions (IBR) of Fc γ RII and Fc ϵ RI, *Immunol. Rev.* 125, 21-35.
- 40 Homsy, J., Meyer, M., Tateno, M., Clarkson, S., Levy, J.A. (1989), The Fc and not CD4 receptor mediates antibody enhancement of HIV infection in human cells, *Science* 244, 1357-1360.
- Hoover, R.G., Lary, C., Page, R., Travis, P., Owens, R., Flick, J., Kornbluth, J., Barlogie, B. (1995), Autoregulatory circuits in myeloma: Tumor cell cytotoxicity mediated by soluble CD16, *J. Clin. Invest.* 95, 241-247.
- 45 Huber, R., Deisenhofer, J., Colman, P.M., Matsushima, M. and Palm, W. (1976), Crystallographic structure studies of an IgG molecule and an Fc fragment, *Nature* 264, 415-420.
- Hulett, M.D., Witort, E., Brinkworth, R.I., McKenzie, I.F.C., Hogarth, P.M. (1994), Identification of the IgG binding site of the human low affinity receptor for IgG Fc γ RII, *J. Biol. Chem.* 269, 15287-15293.
- 50 Hulett, M.D., Witort, E., Brinkworth, R.I., McKenzie, I.F.C., Hogarth, P.M. (1995), Multiple regions of human Fc γ RII (CD32) contribute to the binding of IgG, *J. Biol. Chem.* 270, 21188-21194.
- 55 Ierino, F.L., Powell, M.S., McKenzie, I.F.C., Hogarth, P.M. (1993), Recombinant soluble human Fc γ RII: Production, characterization, and inhibition of the arthus reaction, *J. Exp. Med.* 178, 1617-1628.

- Jancarik, J. and Kim, S.H. (1991), Sparse matrix sampling: A screening method for crystallization of proteins, *J. Appl. Crystallogr.* 24, 409-411.
- 5 Jones, T.A., Zou, J.-Y., Cowan, S.W., Kjeldgaard, M. (1991), Improved methods for building protein models in electron density maps and the location of errors in these models, *Acta crystallogr.* A47, 110-119.
- Khayat, D., Soubrane, C., Andrieu, J.M., Visonneau, S., Eme, D., Tourani, J.M., Beldjord, K., Weil, M., Fernandez, E., Jaquillat, C. (1990), Changes of soluble CD16 levels in serum of HIV patients: Correlation with clinical and biological prognostic factors, *J. Infect. Dis.* 161, 430-435.
- 10 Kraulis, P.J. (1991), MOLSCRIPT: a program to produce both detailed and schematic plots of protein structures, *J. Appl. Cryst.* 24, 946-950.
- Leslie, A.G.W. (1997), Mosflm user guide, mosflm version 5.50, MRC Laboratory of Molecular Biology, Cambridge, UK.
- 15 Lessel, U. and Schomburg, D. (1994), Similarities between protein 3-D structures, *Protein Eng.* 7, 1175-1187.
- Littau, R., Kurane, I. and Ennis, F.A. (1990), Human IgG Fc receptor II mediates antibody-dependent enhancement of dengue virus infection, *J. Immunol.* 144, 3183-3186.
- 20 Lynch, R.G., Hagen, M., Mueller, A., Sandor, M. (1995), Potential role of FcγR in early development of murine lymphoid cells: Evidence for functional interaction between FcγR on pre-thymocytes and an alternative, non-Ig ligand on thymic stromal cells, *Immunol. Lett.* 44, 105-109.
- 25 Mathiot, C., Teillaud, J.L., Elmalek, M., Mosseri, L., Euler-Ziegler, L., Daragon, A., Grosbois, B., Michaux, J.L., Facon, T., Bernard, J.F., Duclos, B., Monconduit, M., Fridman, W.H. (1993), Correlation between serum soluble CD16 (sCD16) levels and disease stage in patients with multiple myeloma, *J. Clin. Immunol.* 13, 41-48.
- Merritt, E.A. and Murphy, M.E.P. (1994), Raster 3D Version 2.0. A program for photorealistic molecular graphics, *Acta Cryst.* D50, 869-873.
- 30 Metzger, H. (1992A), Transmembrane signaling: The joy of aggregation, *J. Immunol.* 149, 1477-1487.
- 35 Metzger, H. (1992B), The receptor with high affinity for Ig E, *Immunol. Rev.* 125, 37-48.
- Müller, S. and Hoover, R.G. (1985), T cells with Fc receptors in myeloma; suppression of growth and secretion of MOPC-315 by T alpha cells, *J. Immunol.* 134, 644-7.
- 40 Nicholls, A., Sharp, K.A., Honig, B. (1991), Protein folding and association: insights from the interfacial and thermodynamic properties of hydrocarbons, *Proteins* 11, 281-296.
- Poo, H., Kraus, J.C., Mayo-Bond, L., Todd, R.F., Petty, H.R. (1995), Interaction of Fcγ receptor IIIB with complement receptor type 3 in fibroblast transfectants: evidence from lateral diffusion and resonance energy transfer studies, *J. Mol. Biol.* 247, 597-603.
- 45 Rappaport, E.F., Cassel, D.L., Walterhouse, D.O., McKenzie, S.E., Surrey, S., Keller, M.A., Schreiber, A.D., Schwartz, E. (1993), A soluble form of the human Fc receptor FcγRIIIa: cloning, transcript analysis and detection. *Exp. Hematol.* 21, 689-696.
- 50 Ravel, K., Castelle, C., Defrance, T., Wild, T.F., Charron, D., Lotteau, V., Rabourdincombe, C. (1997), Measles virus nucleocapsid protein binds to FcγRII and inhibits human B cell antibody production. *J. Exp. Med.* 186, 269-278.
- 55 Roman, S., Moore, J.S., Darby, C., Muller, S., Hoover, R.G. (1988), Modulation of Ig gene expression by Ig binding factors. Suppression of alpha-H chain and lambda-2-L chain mRNA accumulation in MOPC-315 by IgA-binding factor, *J. Immunology* 140, 3622-30.

Sauer-Eriksson, A.E., Kleywegt, G.J., Uhlen, M., Jones, T.A. (1995), Crystal structure of the C2 fragment of streptococcal protein G in complex with the Fc domain of human IgG, *Structure* 3, 265-78.

Sondermann, P., Huber, R., Jacob, U. (1998B), Preparation and crystallization of active soluble human FcγRIIb derived from E.coli, *Protein Structure*, submitted.

Sondermann, P., Kutscher, C., Jacob, U., Frey, J. (1998A), Characterization and crystallization of soluble human Fcγ receptor 11 isoforms produced in insect cells, *Biochemistry*, submitted.

Sondermann, P., Kutscher, C., Jacob, U., Frey, J., Analysis of complexes of IgG and soluble human Fcγ-Receptor II produced in insect cells and its crystallization, submitted.

Tax, W.J.M., Willems, H.W., Reekers, P.P.M., Capel, P.J.A., Koene, R.A.P. (1983), Polymorphism in mitogenic effect of IgG1 monoclonal antibodies against T3 antigen on human T cells, *Nature* 304, 445-447.

Teillaud, J.L., Brunati, S., Elmalek, M., Astier, A., Nicaise, P., Moncuit, J., Mathiot, C., Job-Deslandre, C., Fridman, W.H. (1990), Involvement of FcR + T cells and of IgG-BF in the control of myeloma cells, *Mol. Immunol.* 27, 1209-17.

Turk, D. (1992), Ph.D. Thesis, TU München, Germany.

Ulvestad, E., Matre, R., Tonder, O. (1988), IgG Fc receptors in sera from patients with Rheumatoid Arthritis and Systemic Lupus Erythematosus, *Scand. J. Rheumatol., Suppl.* 75, 203-208.

van de Winkel, J.G.J. and Capel, P.J.A. (1993), Human IgG Fc receptor heterogeneity: Molecular aspects and clinical implications, *Immunol. Today* 14, 215-221.

Varin, N., Sautès, C., Galinha, A., Even, J., Hogarth, P.M., Fridman, W.H. (1989), Recombinant soluble receptors for the Fcγ portion inhibit antibody production in vitro, *Eur. J. Immunol.* 19, 2263-2268.

Yang, Z., Delgado, R., Xu, L., Todd, R.F., Nabel, E.G., Sanchez, A., Nabel, G.J. (1998), Distinct cellular interactions of secreted and transmembrane Ebola virus glycoproteins, *Science* 279, 983-984.

Zhou, M.-J., Todd, R.F., van de Winkel, J.G.J., Petty, H.R. (1993), Cocapping of the leukoadhesin molecules complement receptor type 3 and lymphocyte function-associated antigen-1 with Fcγ receptor III on human neutrophils. Possible role of lectin-like interactions, *J. Immunol.* 150, 3030-3041.

SEQUENZPROTOKOLL

5 <110> Max-Planck-Gesellschaft zur Förderung der Wissensc
 <120> Recombinant soluble Fc receptors
 <130> recombinant soluble fc receptors
 10 <140> 98 122 969.3
 <141> 1998-12-03
 <160> 14
 15 <170> PatentIn Ver. 2.1
 <210> 1
 <211> 1321
 20 <212> DNA
 <213> Homo sapiens
 <220>
 25 <221> CDS
 <222> (37)..(1158)
 <400> 1
 gacagatttc actgctcca ccagcttga gacaac atg tgg ttc ttg aca act 54
 30 Met Trp Phe Leu Thr Thr
 1 5
 ctg ctc ctt tgg gtt cca gtt gat ggg caa gtg gac acc aca aag gca 102
 35 Leu Leu Leu Trp Val Pro Val Asp Gly Gln Val Asp Thr Thr Lys Ala
 10 15 20
 gtg atc tct ttg cag cct cca tgg gtc agc gtg ttc caa gag gaa acc 150
 Val Ile Ser Leu Gln Pro Pro Trp Val Ser Val Phe Gln Glu Glu Thr
 25 30 35
 gta acc ttg cac tgt gag gtg ctc cat ctg cct ggg agc agc tct aca 198
 Val Thr Leu His Cys Glu Val Leu His Leu Pro Gly Ser Ser Ser Thr
 40 45 50
 cag tgg ttt ctc aat ggc aca gcc act cag acc tgg acc ccc agc tac 246
 Gln Trp Phe Leu Asn Gly Thr Ala Thr Gln Thr Ser Thr Pro Ser Tyr
 55 60 65 70
 aga atc acc tct gcc agt gtc aat gac agt ggt gaa tac agg tgc cag 294
 Arg Ile Thr Ser Ala Ser Val Asn Asp Ser Gly Glu Tyr Arg Cys Gln
 75 80 85

55

EP 1 006 183 A1

5	aga ggt ctc tca ggg cga agt gac ccc ata cag ctg gaa atc cac aga	342
	Arg Gly Leu Ser Gly Arg Ser Asp Pro Ile Gln Leu Glu Ile His Arg	
	90 95 100	
10	ggc tgg cta cta ctg cag gtc tcc agc aga gtc ttc acg gaa gga gaa	390
	Gly Trp Leu Leu Leu Gln Val Ser Ser Arg Val Phe Thr Glu Gly Glu	
	105 110 115	
15	cct ctg gcc ttg agg tgt cat gcg tgg aag gat aag ctg gtg tac aat	438
	Pro Leu Ala Leu Arg Cys His Ala Trp Lys Asp Lys Leu Val Tyr Asn	
	120 125 130	
20	gtg ctt tac tat cga aat ggc aaa gcc ttt aag ttt ttc cac tgg aat	486
	Val Leu Tyr Tyr Arg Asn Gly Lys Ala Phe Lys Phe Phe His Trp Asn	
	135 140 145 150	
25	tct aac ctc acc att ctg aaa acc aac ata agt cac aat ggc acc tac	534
	Ser Asn Leu Thr Ile Leu Lys Thr Asn Ile Ser His Asn Gly Thr Tyr	
	155 160 165	
30	cat tgc tca ggc atg gga aag cat cgc tac aca tca gca gga ata tct	582
	His Cys Ser Gly Met Gly Lys His Arg Tyr Thr Ser Ala Gly Ile Ser	
	170 175 180	
35	gtc act gtg aaa gag cta ttt cca gct cca gtg ctg aat gca tct gtg	630
	Val Thr Val Lys Glu Leu Phe Pro Ala Pro Val Leu Asn Ala Ser Val	
	185 190 195	
40	aca tcc cca ctc ctg gag ggg aat ctg gtc acc ctg agc tgt gaa aca	678
	Thr Ser Pro Leu Leu Glu Gly Asn Leu Val Thr Leu Ser Cys Glu Thr	
	200 205 210	
45	aag ttg ctc ttg cag agg cct ggt ttg cag ctt tac ttc tcc ttc tac	726
	Lys Leu Leu Leu Gln Arg Pro Gly Leu Gln Leu Tyr Phe Ser Phe Tyr	
	215 220 225 230	
50	atg ggc agc aag acc ctg cga ggc agg aac aca tcc tct gaa tac caa	774
	Met Gly Ser Lys Thr Leu Arg Gly Arg Asn Thr Ser Ser Glu Tyr Gln	
	235 240 245	
55	ata cta act gct aga aga gaa gac tct ggg tta tac tgg tgc gag gct	822
	Ile Leu Thr Ala Arg Arg Glu Asp Ser Gly Leu Tyr Trp Cys Glu Ala	
	250 255 260	
60	gcc aca gag gat gga aat gtc ctt aag cgc agc cct gag ttg gag ctt	870
	Ala Thr Glu Asp Gly Asn Val Leu Lys Arg Ser Pro Glu Leu Glu Leu	
	265 270 275	

EP 1 006 183 A1

5 caa gtg ctt ggc ctc cag tta cca act cct gtc tgg ttt cat gtc ctt 918
 Gln Val Leu Gly Leu Gln Leu Pro Thr Pro Val Trp Phe His Val Leu
 280 285 290

10 ttc tat ctg gca gtg gga ata atg ttt tta gtg aac act gtt ctc tgg 966
 Phe Tyr Leu Ala Val Gly Ile Met Phe Leu Val Asn Thr Val Leu Trp
 295 300 305 310

15 gtg aca ata cgt aaa gaa ctg aaa aga aag aaa aag tgg gat tta gaa 1014
 Val Thr Ile Arg Lys Glu Leu Lys Arg Lys Lys Lys Trp Asp Leu Glu
 315 320 325

20 atc tct ttg gat tct ggt cat gag aag aag gta act tcc agc ctt caa 1062
 Ile Ser Leu Asp Ser Gly His Glu Lys Lys Val Thr Ser Ser Leu Gln
 330 335 340

25 gaa gac aga cat tta gaa gaa gag ctg aaa tgt cag gaa caa aaa gaa 1110
 Glu Asp Arg His Leu Glu Glu Glu Leu Lys Cys Gln Glu Gln Lys Glu
 345 350 355

30 gaa cag ctg cag gaa ggg gtg cac cgg aag gag ccc cag ggg gcc acg 1158
 Glu Gln Leu Gln Glu Gly Val His Arg Lys Glu Pro Gln Gly Ala Thr
 360 365 370

35 tagcagcggc tcagtgggtg gccatcgatc tggaccgtcc cctgcccact tgctccccgt 1218
 gagcactgcg tacaaacatc caaaagtcca acaacaccag aactgtgtgt ctcattggtat 1278
 gtaactctta aagcaaataa atgaactgac ttcaaaaaaa aaa 1321

40 <210> 2
 <211> 374
 <212> PRT
 <213> Homo sapiens

45 <400> 2
 Met Trp Phe Leu Thr Thr Leu Leu Leu Trp Val Pro Val Asp Gly Gln
 1 5 10 15

50 Val Asp Thr Thr Lys Ala Val Ile Ser Leu Gln Pro Pro Trp Val Ser
 20 25 30

55 Val Phe Gln Glu Glu Thr Val Thr Leu His Cys Glu Val Leu His Leu
 35 40 45

 Pro Gly Ser Ser Ser Thr Gln Trp Phe Leu Asn Gly Thr Ala Thr Gln

EP 1 006 183 A1

	50	55	60
5	Thr Ser Thr Pro Ser Tyr Arg Ile Thr Ser Ala Ser Val Asn Asp Ser 65 70 75 80		
	Gly Glu Tyr Arg Cys Gln Arg Gly Leu Ser Gly Arg Ser Asp Pro Ile 85 90 95		
10	Gln Leu Glu Ile His Arg Gly Trp Leu Leu Leu Gln Val Ser Ser Arg 100 105 110		
	Val Phe Thr Glu Gly Glu Pro Leu Ala Leu Arg Cys His Ala Trp Lys 115 120 125		
15	Asp Lys Leu Val Tyr Asn Val Leu Tyr Tyr Arg Asn Gly Lys Ala Phe 130 135 140		
20	Lys Phe Phe His Trp Asn Ser Asn Leu Thr Ile Leu Lys Thr Asn Ile 145 150 155 160		
	Ser His Asn Gly Thr Tyr His Cys Ser Gly Met Gly Lys His Arg Tyr 165 170 175		
25	Thr Ser Ala Gly Ile Ser Val Thr Val Lys Glu Leu Phe Pro Ala Pro 180 185 190		
30	Val Leu Asn Ala Ser Val Thr Ser Pro Leu Leu Glu Gly Asn Leu Val 195 200 205		
	Thr Leu Ser Cys Glu Thr Lys Leu Leu Leu Gln Arg Pro Gly Leu Gln 210 215 220		
35	Leu Tyr Phe Ser Phe Tyr Met Gly Ser Lys Thr Leu Arg Gly Arg Asn 225 230 235 240		
40	Thr Ser Ser Glu Tyr Gln Ile Leu Thr Ala Arg Arg Glu Asp Ser Gly 245 250 255		
	Leu Tyr Trp Cys Glu Ala Ala Thr Glu Asp Gly Asn Val Leu Lys Arg 260 265 270		
45	Ser Pro Glu Leu Glu Leu Gln Val Leu Gly Leu Gln Leu Pro Thr Pro 275 280 285		
50	Val Trp Phe His Val Leu Phe Tyr Leu Ala Val Gly Ile Met Phe Leu 290 295 300		
	Val Asn Thr Val Leu Trp Val Thr Ile Arg Lys Glu Leu Lys Arg Lys		

55

EP 1 006 183 A1

	305		310		315		320
5	Lys Lys Trp Asp Leu Glu Ile Ser Leu Asp Ser Gly His Glu Lys Lys	325		330		335	
	Val Thr Ser Ser Leu Gln Glu Asp Arg His Leu Glu Glu Glu Leu Lys	340		345		350	
10	Cys Gln Glu Gln Lys Glu Glu Gln Leu Gln Glu Gly Val His Arg Lys	355		360		365	
15	Glu Pro Gln Gly Ala Thr	370					
	<210> 3						
20	<211> 1419						
	<212> DNA						
	<213> Homo sapiens						
	<220>						
25	<221> CDS						
	<222> (45)..(917)						
	<400> 3						
30	ggctgtgact gctgtgctct gggcgccact cgctccaggg agtg atg gga atc ctg	56					
					Met Gly Ile Leu		
					1		
	tca ttt tta cct gtc ctt gcc act gag agt gac tgg gct gac tgc aag	104					
35	Ser Phe Leu Pro Val Leu Ala Thr Glu Ser Asp Trp Ala Asp Cys Lys						
	5 10 15 20						
	tcc ccc cag cct tgg ggt cat atg ctt ctg tgg aca gct gtg cta ttc	152					
40	Ser Pro Gln Pro Trp Gly His Met Leu Leu Trp Thr Ala Val Leu Phe						
	25 30 35						
	ctg gct cct gtt gct ggg aca cct gca gct ccc cca aag gct gtg ctg	200					
45	Leu Ala Pro Val Ala Gly Thr Pro Ala Ala Pro Pro Lys Ala Val Leu						
	40 45 50						
	aaa ctc gag ccc cag tgg atc aac gtg ctc cag gag gac tct gtg act	248					
50	Lys Leu Glu Pro Gln Trp Ile Asn Val Leu Gln Glu Asp Ser Val Thr						
	55 60 65						
	ctg aca tgc cgg ggg act cac agc cct gag agc gac tcc att cag tgg	296					
55	Leu Thr Cys Arg Gly Thr His Ser Pro Glu Ser Asp Ser Ile Gln Trp						
	70 75 80						

55

EP 1 006 183 A1

atg cac ccg gat gct ctg gaa gag cct gat gac cag aac cgt att 917
 Met His Pro Asp Ala Leu Glu Glu Pro Asp Asp Gln Asn Arg Ile
 5 280 285 290

 tagttcccat tgtcttgcac tgggatttga gaagaaatca gagagggaag atctggtatt 977
 tcttggccta aattccccctt ggggaggaca gggagatgct gcagttccaa aagagaaggt 1037
 10 tttctccaga gtcactctacc tgagtectga agctccctgt cctgaaagcc acagacaata 1097
 tgggtcccaaa tgcccgactg caocttctgt gcttcagctc ttottgacat caaggctctt 1157
 15 ccgttccaca tccacacagc caatccaatt aatcaaacca ctgttattaa cagataatag 1217
 caacttggga aatgcttatg ttacaggtta cgtgagaaca atcatgtaaa tctatatgat 1277
 20 ttcagaaatg ttaaaataga ctaacctcta ccagcacatt aaaagtgatt gtttctgggt 1337
 gataaaatta ttgatgattt ttattttctt tatttttcta taaagatcat atattacttt 1397
 25 tataataaaa cattataaaa ac 1419

 <210> 4
 <211> 291
 30 <212> PRT
 <213> Homo sapiens

 <400> 4
 35 Met Gly Ile Leu Ser Phe Leu Pro Val Leu Ala Thr Glu Ser Asp Trp
 1 5 10 15
 Ala Asp Cys Lys Ser Pro Gln Pro Trp Gly His Met Leu Leu Trp Thr
 40 20 25 30
 Ala Val Leu Phe Leu Ala Pro Val Ala Gly Thr Pro Ala Ala Pro Pro
 35 40 45
 45 Lys Ala Val Leu Lys Leu Glu Pro Gln Trp Ile Asn Val Leu Gln Glu
 50 55 60
 Asp Ser Val Thr Leu Thr Cys Arg Gly Thr His Ser Pro Glu Ser Asp
 50 65 70 75 80
 Ser Ile Gln Trp Phe His Asn Gly Asn Leu Ile Pro Thr His Thr Gln
 85 90 95
 55

EP 1 006 183 A1

5 Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn Asp Ser Gly Glu Tyr Thr
 100 105 110
 6 Cys Gln Thr Gly Gln Thr Ser Leu Ser Asp Pro Val His Leu Thr Val
 115 120 125
 10 Leu Ser Glu Trp Leu Val Leu Gln Thr Pro His Leu Glu Phe Gln Glu
 130 135 140
 Gly Glu Thr Ile Val Leu Arg Cys His Ser Trp Lys Asp Lys Pro Leu
 145 150 155 160
 15 Val Lys Val Thr Phe Phe Gln Asn Gly Lys Ser Lys Lys Phe Ser Arg
 165 170 175
 20 Ser Asp Pro Asn Phe Ser Ile Pro Gln Ala Asn His Ser His Ser Gly
 180 185 190
 Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr Thr Leu Tyr Ser Ser Lys
 195 200 205
 25 Pro Val Thr Ile Thr Val Gln Ala Pro Ser Ser Ser Pro Met Gly Ile
 210 215 220
 30 Ile Val Ala Val Val Thr Gly Ile Ala Val Ala Ala Ile Val Ala Ala
 225 230 235 240
 Val Val Ala Leu Ile Tyr Cys Arg Lys Lys Arg Ile Ser Ala Asn Pro
 245 250 255
 35 Thr Asn Pro Asp Glu Ala Asp Lys Val Gly Ala Glu Asn Thr Ile Thr
 260 265 270
 Tyr Ser Leu Leu Met His Pro Asp Ala Leu Glu Glu Pro Asp Asp Gln
 275 280 285
 40 Asn Arg Ile
 290
 45 <210> 5
 <211> 2359
 <212> DNA
 <213> Homo sapiens
 50 <220>
 <221> CDS
 <222> (6) .. (938)
 55

EP 1 006 183 A1

<400> 5
 cccaa atg tct cag aat gta tgt ccc aga aac ctg tgg ctg ctt caa cca 50
 Met Ser Gln Asn Val Cys Pro Arg Asn Leu Trp Leu Leu Gln Pro
 1 5 10 15

ttg aca gtt ttg ctg ctg ctg gct tct gca gac agt caa gct gca gct 98
 Leu Thr Val Leu Leu Leu Leu Ala Ser Ala Asp Ser Gln Ala Ala Ala
 20 25 30

ccc cca aag gct gtg ctg aaa ctt gag ccc ccg tgg atc aac gtg ctc 146
 Pro Pro Lys Ala Val Leu Lys Leu Glu Pro Pro Trp Ile Asn Val Leu
 35 40 45

cag gag gac tct gtg act ctg aca tgc cag ggg gct cgc agc cct gag 194
 Gln Glu Asp Ser Val Thr Leu Thr Cys Gln Gly Ala Arg Ser Pro Glu
 50 55 60

agc gac tcc att cag tgg ttc cac aat ggg aat ctc att ccc acc cac 242
 Ser Asp Ser Ile Gln Trp Phe His Asn Gly Asn Leu Ile Pro Thr His
 65 70 75

acg cag ccc agc tac agg ttc aag gcc aac aac aat gac agc ggg gag 290
 Thr Gln Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn Asp Ser Gly Glu
 80 85 90 95

tac acg tgc cag act ggc cag acc agc ctc agc gac cct gtg cat ctg 338
 Tyr Thr Cys Gln Thr Gly Gln Thr Ser Leu Ser Asp Pro Val His Leu
 100 105 110

act gtg ctt tcc gaa tgg ctg gtg ctc cag acc cct cac ctg gag ttc 386
 Thr Val Leu Ser Glu Trp Leu Val Leu Gln Thr Pro His Leu Glu Phe
 115 120 125

cag gag gga gaa acc atc atg ctg agg tgc cac agc tgg aag gac aag 434
 Gln Glu Gly Glu Thr Ile Met Leu Arg Cys His Ser Trp Lys Asp Lys
 130 135 140

cct ctg gtc aag gtc aca ttc ttc cag aat gga aaa tcc cag aaa ttc 482
 Pro Leu Val Lys Val Thr Phe Phe Gln Asn Gly Lys Ser Gln Lys Phe
 145 150 155

tcc cgt ttg gat ccc acc ttc tcc atc cca caa gca aac cac agt cac 530
 Ser Arg Leu Asp Pro Thr Phe Ser Ile Pro Gln Ala Asn His Ser His
 160 165 170 175

agt ggt gat tac cac tgc aca gga aac ata ggc tac acg ctg ttc tca 578
 Ser Gly Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr Thr Leu Phe Ser

EP 1 006 183 A1

	180	185	190	
5	tcc aag cct gtg acc atc act gtc caa gtg ccc agc atg ggc agc tct			626
	Ser Lys Pro Val Thr Ile Thr Val Gln Val Pro Ser Met Gly Ser Ser			
	195	200	205	
10	tca cca atg ggg atc att gtg gct gtg gtc att gcg act gct gta gca			674
	Ser Pro Met Gly Ile Ile Val Ala Val Val Ile Ala Thr Ala Val Ala			
	210	215	220	
15	gcc att gtt gct gct gta gtg gcc ttg atc tac tgc agg aaa aag cgg			722
	Ala Ile Val Ala Ala Val Val Ala Leu Ile Tyr Cys Arg Lys Lys Arg			
	225	230	235	
20	att tca gcc aat tcc act gat cct gtg aag gct gcc caa ttt gag cca			770
	Ile Ser Ala Asn Ser Thr Asp Pro Val Lys Ala Ala Gln Phe Glu Pro			
	240	245	250	255
25	cct gga cgt caa atg att gcc atc aga aag aga caa ctt gaa gaa acc			818
	Pro Gly Arg Gln Met Ile Ala Ile Arg Lys Arg Gln Leu Glu Glu Thr			
	260	265	270	
30	aac aat gac tat gaa aca gct gac ggc ggc tac atg act ctg aac ccc			866
	Asn Asn Asp Tyr Glu Thr Ala Asp Gly Gly Tyr Met Thr Leu Asn Pro			
	275	280	285	
35	agg gca cct act gac gat gat aaa aac atc tac ctg act ctt cct ccc			914
	Arg Ala Pro Thr Asp Asp Asp Lys Asn Ile Tyr Leu Thr Leu Pro Pro			
	290	295	300	
40	aac gac cat gtc aac agt aat aac taaagagtaa cgttatgcc a tgtggtcata			968
	Asn Asp His Val Asn Ser Asn Asn			
	305	310		
45	ctctcagcctt gctgatggat gacaaaaaga ggggaattgt taaaggaaaa tttaaatgga			1028
	gactggaaaa atcctgagca aacaaaacca cctggccctt agaaatagct ttaactttgc			1088
	ttaaactaca aacacaagca aaacttcacg gggtcatact acatacaagc ataagcaaaa			1148
	cttaacttgg atcatttctg gttaatgctt atgttagaaa taagacaacc ccagccaatc			1208
	acaagcagcc tactaacata taattagggt actagggact ttctaagaag atacctaccc			1268
50	ccaaaaaaca attatgtaat tgaaaaccaa ccgattgcct ttattttgct tccacatttt			1328
	cccaataaat acttgccctgt gacattttgc cactggaaca ctaaacttca tgaattgcgc			1388
55				

EP 1 006 183 A1

ctcagatttt tcccttaaca tctttttttt ttttgacaga gtctcaatct gttaccag 1448
5 ctggagtga gtggtgctat cttggtcac tgcaaaccg cctcccaggt ttaagcgatt 1508
cttatgcctc agcctcccag tagctgggat tagaggcatg tgccatcata cccagcta 1568
10 ttttgtattt tttattttt atttttagta gagacagggg ttcgcaatgt tggccaggcc 1628
gatctcgaac ttctggcctc tagcgtctg cccgcctcgg cctcccaaag tgctgggatg 1688
accgcatcag ccccaatgtc cagcctcttt aacatcttct ttcctatgcc ctctctgtgg 1748
15 atccctactg ctggtttctg cctctcccat gctgagaaca aaatcaccta ttcactgctt 1808
atgcagtcgg aagctccaga agaacaaaga gcccaattac cagaaccaca ttaagtctcc 1868
20 attgttttgc cttgggattt gagaagagaa ttagagaggt gaggatctgg tatttcctgg 1928
actaaattcc cttggggaag acgaagggat gctgcagttc caaaagagaa ggactcttcc 1988
agagtcattc acctgagtc caaagctccc tgtcctgaaa gccacagaca atatgggtccc 2048
25 aaatgactga ctgcacctc tgtgcctcag ccgttcttga catcaagaat cttctgttcc 2108
acatccacac agccaatata attagtcaaa ccactgttat taacagatgt agcaacatga 2168
30 gaaacgctta tgttacaggt tacatgagag caatcatgta agtctatatg acttcagaaa 2228
tgttaaaata gactaacctc taacaacaaa ttaaaagtga ttgtttcaag gtgatgcaat 2288
tattgatgac ctattttatt tttctataat gatcatatat taccttttga ataaaacatt 2348
35 ataaccaaaa c 2359

40 <210> 6
<211> 311
<212> PRT
<213> Homo sapiens

45 <400> 6
Met Ser Gln Asn Val Cys Pro Arg Asn Leu Trp Leu Leu Gln Pro Leu
1 5 10 15
50 Thr Val Leu Leu Leu Leu Ala Ser Ala Asp Ser Gln Ala Ala Ala Pro
20 25 30
Pro Lys Ala Val Leu Lys Leu Glu Pro Pro Trp Ile Asn Val Leu Gln

55

EP 1 006 183 A1

	35	40	45
5	Glu Asp Ser Val Thr Leu Thr Cys Gln Gly Ala Arg Ser Pro Glu Ser 50	55	60
	Asp Ser Ile Gln Trp Phe His Asn Gly Asn Leu Ile Pro Thr His Thr 65	70	75 80
10	Gln Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn Asp Ser Gly Glu Tyr 85	90	95
	Thr Cys Gln Thr Gly Gln Thr Ser Leu Ser Asp Pro Val His Leu Thr 100	105	110
15	Val Leu Ser Glu Trp Leu Val Leu Gln Thr Pro His Leu Glu Phe Gln 115	120	125
	Glu Gly Glu Thr Ile Met Leu Arg Cys His Ser Trp Lys Asp Lys Pro 130	135	140
20	Leu Val Lys Val Thr Phe Phe Gln Asn Gly Lys Ser Gln Lys Phe Ser 145	150	155 160
25	Arg Leu Asp Pro Thr Phe Ser Ile Pro Gln Ala Asn His Ser His Ser 165	170	175
	Gly Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr Thr Leu Phe Ser Ser 180	185	190
30	Lys Pro Val Thr Ile Thr Val Gln Val Pro Ser Met Gly Ser Ser Ser 195	200	205
35	Pro Met Gly Ile Ile Val Ala Val Val Ile Ala Thr Ala Val Ala Ala 210	215	220
	Ile Val Ala Ala Val Val Ala Leu Ile Tyr Cys Arg Lys Lys Arg Ile 225	230	235 240
40	Ser Ala Asn Ser Thr Asp Pro Val Lys Ala Ala Gln Phe Glu Pro Pro 245	250	255
	Gly Arg Gln Met Ile Ala Ile Arg Lys Arg Gln Leu Glu Glu Thr Asn 260	265	270
45	Asn Asp Tyr Glu Thr Ala Asp Gly Gly Tyr Met Thr Leu Asn Pro Arg 275	280	285
50	Ala Pro Thr Asp Asp Asp Lys Asn Ile Tyr Leu Thr Leu Pro Pro Asn		
55			

EP 1 006 183 A1

	290	295	300	
5	Asp His Val Asn Ser Asn Asn			
	305	310		
10	<210> 7 <211> 887 <212> DNA <213> Homo sapiens			
15	<220> <221> CDS <222> (34)..(732)			
20	<400> 7 tctttggtga cttgtccact ccagtggtgc atc atg tgg cag ctg ctc ctc cca 54 <div style="text-align: center;">Met Trp Gln Leu Leu Leu Pro</div> <div style="display: flex; justify-content: space-around; width: 100%;"> 1 5 </div>			
25	act gct ctg cta ctt cta gtt tca gct ggc atg cgg act gaa gat ctc 102 Thr Ala Leu Leu Leu Leu Val Ser Ala Gly Met Arg Thr Glu Asp Leu <div style="display: flex; justify-content: space-around; width: 100%;"> 10 15 20 </div>			
30	cca aag gct gtg gtg ttc ctg gag cct caa tgg tac agc gtg ctt gag 150 Pro Lys Ala Val Val Phe Leu Glu Pro Gln Trp Tyr Ser Val Leu Glu <div style="display: flex; justify-content: space-around; width: 100%;"> 25 30 35 </div>			
35	aag gac agt gtg act ctg aag tgc cag gga gcc tac tcc cct gag gac 198 Lys Asp Ser Val Thr Leu Lys Cys Gln Gly Ala Tyr Ser Pro Glu Asp <div style="display: flex; justify-content: space-around; width: 100%;"> 40 45 50 55 </div>			
40	aat tcc aca cag tgg ttt cac aat gag agc ctc atc tca agc cag gcc 246 Asn Ser Thr Gln Trp Phe His Asn Glu Ser Leu Ile Ser Ser Gln Ala <div style="display: flex; justify-content: space-around; width: 100%;"> 60 65 70 </div>			
45	tcg agc tac ttc att gac gct gcc aca gtc aac gac agt gga gag tac 294 Ser Ser Tyr Phe Ile Asp Ala Ala Thr Val Asn Asp Ser Gly Glu Tyr <div style="display: flex; justify-content: space-around; width: 100%;"> 75 80 85 </div>			
50	agg tgc cag aca aac ctc tcc acc ctc agt gac ccg gtg cag cta gaa 342 Arg Cys Gln Thr Asn Leu Ser Thr Leu Ser Asp Pro Val Gln Leu Glu <div style="display: flex; justify-content: space-around; width: 100%;"> 90 95 100 </div>			
55	gtc cat atc ggc tgg ctg ttg ctc cag gcc cct cgg tgg gtg ttc aag 390 Val His Ile Gly Trp Leu Leu Leu Gln Ala Pro Arg Trp Val Phe Lys <div style="display: flex; justify-content: space-around; width: 100%;"> 105 110 115 </div>			

EP 1 006 183 A1

5 gag gaa gac cct att cac ctg agg tgt cac agc tgg aag aac act gct 438
 Glu Glu Asp Pro Ile His Leu Arg Cys His Ser Trp Lys Asn Thr Ala
 120 125 130 135

10 ctg cat aag gtc aca tat tta cag aat ggc aaa gac agg aag tat ttt 486
 Leu His Lys Val Thr Tyr Leu Gln Asn Gly Lys Asp Arg Lys Tyr Phe
 140 145 150

15 cat cat aat tct gac ttc cac att cca aaa gcc aca ctc aaa gat agc 534
 His His Asn Ser Asp Phe His Ile Pro Lys Ala Thr Leu Lys Asp Ser
 155 160 165

20 ggc tcc tac ttc tgc agg ggg ctt gtt ggg agt aaa aat gtg tct tca 582
 Gly Ser Tyr Phe Cys Arg Gly Leu Val Gly Ser Lys Asn Val Ser Ser
 170 175 180

25 gag act gtg aac atc acc atc act caa ggt ttg gca gtg tca acc atc 630
 Glu Thr Val Asn Ile Thr Ile Thr Gln Gly Leu Ala Val Ser Thr Ile
 185 190 195

30 tca tca ttc tct cca cct ggg tac caa gtc tct ttc tgc ttg gtg atg 678
 Ser Ser Phe Ser Pro Pro Gly Tyr Gln Val Ser Phe Cys Leu Val Met
 200 205 210 215

35 gta ctc ett ttt gca gtg gac aca gga cta tat ttc tct gtg aag aca 726
 Val Leu Leu Phe Ala Val Asp Thr Gly Leu Tyr Phe Ser Val Lys Thr
 220 225 230

40 aac att tgaagctcaa caagagactg gaaggacat aaacttaaat ggagaaagga 782
 Asn Ile

45 ccctcaagac aaatgacccc catcccatgg gagtaataag agcagtggca gcagcatctc 842

50 tgaacatttc tctggatttg caaccccatc atcctcaggc ctctc 887

<210> 8
 <211> 233
 <212> PRT
 <213> Homo sapiens

55 <400> 8
 Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu Leu Val Ser Ala
 1 5 10 15
 Gly Met Arg Thr Glu Asp Leu Pro Lys Ala Val Val Phe Leu Glu Pro
 20 25 30

EP 1 006 183 A1

5 Gln Trp Tyr Ser Val Leu Glu Lys Asp Ser Val Thr Leu Lys Cys Gln
 35 40 45
 10 Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Gln Trp Phe His Asn Glu
 50 55 60
 15 Ser Leu Ile Ser Ser Gln Ala Ser Ser Tyr Phe Ile Asp Ala Ala Thr
 65 70 75 80
 20 Val Asn Asp Ser Gly Glu Tyr Arg Cys Gln Thr Asn Leu Ser Thr Leu
 85 90 95
 25 Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp Leu Leu Leu Gln
 100 105 110
 30 Ala Pro Arg Trp Val Phe Lys Glu Glu Asp Pro Ile His Leu Arg Cys
 115 120 125
 35 His Ser Trp Lys Asn Thr Ala Leu His Lys Val Thr Tyr Leu Gln Asn
 130 135 140
 40 Gly Lys Asp Arg Lys Tyr Phe His His Asn Ser Asp Phe His Ile Pro
 145 150 155 160
 45 Lys Ala Thr Leu Lys Asp Ser Gly Ser Tyr Phe Cys Arg Gly Leu Val
 165 170 175
 50 Gly Ser Lys Asn Val Ser Ser Glu Thr Val Asn Ile Thr Ile Thr Gln
 180 185 190
 55 Gly Leu Ala Val Ser Thr Ile Ser Ser Phe Ser Pro Pro Gly Tyr Gln
 195 200 205
 60 Val Ser Phe Cys Leu Val Met Val Leu Leu Phe Ala Val Asp Thr Gly
 210 215 220
 65 Leu Tyr Phe Ser Val Lys Thr Asn Ile
 225 230
 70 <210> 9
 <211> 1068
 <212> DNA
 <213> Homo sapiens
 75 <220>
 <221> CDS
 <222> (31)..(810)

EP 1 006 183 A1

<400> 9
 agatctcagc acagtaagca ccaggagtc atg aag aag atg gct cct gcc atg 54
 5 Met Lys Lys Met Ala Pro Ala Met
 1 5

gaa tcc cct act cta ctg tgt gta gcc tta ctg ttc ttc gct cca gat 102
 10 Glu Ser Pro Thr Leu Leu Cys Val Ala Leu Leu Phe Phe Ala Pro Asp
 10 15 20

ggc gtg tta gca gtc cct cag aaa cct aag gtc tcc ttg aac cct cca 150
 15 Gly Val Leu Ala Val Pro Gln Lys Pro Lys Val Ser Leu Asn Pro Pro
 25 30 35 40

tgg aat aga ata ttt aaa gga gag aat gtg act ctt aca tgt aat ggg 198
 20 Trp Asn Arg Ile Phe Lys Gly Glu Asn Val Thr Leu Thr Cys Asn Gly
 45 50 55

aac aat ttc ttt gaa gtc agt tcc acc aaa tgg ttc cac aat ggc agc 246
 Asn Asn Phe Phe Glu Val Ser Ser Thr Lys Trp Phe His Asn Gly Ser
 60 65 70

ctt tca gaa gag aca aat tca agt ttg aat att gtg aat gcc aaa ttt 294
 25 Leu Ser Glu Glu Thr Asn Ser Ser Leu Asn Ile Val Asn Ala Lys Phe
 75 80 85

gaa gac agt gga gaa tac aaa tgt cag cac caa caa gtt aat gag agt 342
 30 Glu Asp Ser Gly Glu Tyr Lys Cys Gln His Gln Gln Val Asn Glu Ser
 90 95 100

gaa cct gtg tac ctg gaa gtc ttc agt gac tgg ctg ctc ctt cag gcc 390
 35 Glu Pro Val Tyr Leu Glu Val Phe Ser Asp Trp Leu Leu Leu Gln Ala
 105 110 115 120

tct gct gag gtg gtg atg gag ggc cag ccc ctc ttc ctc agg tgc cat 438
 40 Ser Ala Glu Val Val Met Glu Gly Gln Pro Leu Phe Leu Arg Cys His
 125 130 135

ggc tgg agg aac tgg gat gtg tac aag gtg atc tat tat aag gat ggt 486
 45 Gly Trp Arg Asn Trp Asp Val Tyr Lys Val Ile Tyr Tyr Lys Asp Gly
 140 145 150

gaa gct ctc aag tac tgg tat gag aac cac aac atc tcc att aca aat 534
 Glu Ala Leu Lys Tyr Trp Tyr Glu Asn His Asn Ile Ser Ile Thr Asn
 155 160 165

gcc aca gtt gaa gac agt gga acc tac tac tgt acg ggc aaa gtg tgg 582
 50 Ala Thr Val Glu Asp Ser Gly Thr Tyr Tyr Cys Thr Gly Lys Val Trp
 55

EP 1 006 183 A1

	170	175	180	
5	cag ctg gac tat gag tct gag ccc ctc aac att act gta ata aaa gct			630
	Gln Leu Asp Tyr Glu Ser Glu Pro Leu Asn Ile Thr Val Ile Lys Ala			
	185	190	195	200
10	ccg cgt gag aag tac tgg cta caa ttt ttt atc cca ttg ttg gtg gtg			678
	Pro Arg Glu Lys Tyr Trp Leu Gln Phe Phe Ile Pro Leu Leu Val Val			
	205	210	215	
15	att ctg ttt gct gtg gac aca gga tta ttt atc tca act cag cag cag			726
	Ile Leu Phe Ala Val Asp Thr Gly Leu Phe Ile Ser Thr Gln Gln Gln			
	220	225	230	
20	gtc aca ttt ctc ttg aag att aag aga acc agg aaa ggc ttc aga ctt			774
	Val Thr Phe Leu Leu Lys Ile Lys Arg Thr Arg Lys Gly Phe Arg Leu			
	235	240	245	
25	ctg aac cca cat cct aag cca aac ccc aaa aac aac tgatataatt			820
	Leu Asn Pro His Pro Lys Pro Asn Pro Lys Asn Asn			
	250	255	260	
30	aactcaagaa atatttgcaa cattagtgtt tttccagcat cagcaattgc tactcaattg			880
	tcaaacacag cttgcaatat acatagaaac gtctgtgctc aaggatttat agaaatgctt			940
	cattaaactg agtgaaactg attaagtggc atgtaatagt aagtgtctcaa ttaacattgg			1000
	ttgaataaat gagagaatga atagattcat ttattagcat ttgtaaaaga gatgttcaat			1060
35	ttagatct			1068
40	<210> 10			
	<211> 260			
	<212> PRT			
	<213> Homo sapiens			
45	<400> 10			
	Met Lys Lys Met Ala Pro Ala Met Glu Ser Pro Thr Leu Leu Cys Val			
	1 5 10 15			
50	Ala Leu Leu Phe Phe Ala Pro Asp Gly Val Leu Ala Val Pro Gln Lys			
	20 25 30			
	Pro Lys Val Ser Leu Asn Pro Pro Trp Asn Arg Ile Phe Lys Gly Glu			
	35 40 45			
55				

EP 1 006 183 A1

5 Asn Val Thr Leu Thr Cys Asn Gly Asn Asn Phe Phe Glu Val Ser Ser
 50 55 60
 10 Thr Lys Trp Phe His Asn Gly Ser Leu Ser Glu Glu Thr Asn Ser Ser
 65 70 75 80
 15 Leu Asn Ile Val Asn Ala Lys Phe Glu Asp Ser Gly Glu Tyr Lys Cys
 85 90 95
 20 Gln His Gln Gln Val Asn Glu Ser Glu Pro Val Tyr Leu Glu Val Phe
 100 105 110
 25 Ser Asp Trp Leu Leu Leu Gln Ala Ser Ala Glu Val Val Met Glu Gly
 115 120 125
 30 Gln Pro Leu Phe Leu Arg Cys His Gly Trp Arg Asn Trp Asp Val Tyr
 130 135 140
 35 Lys Val Ile Tyr Tyr Lys Asp Gly Glu Ala Leu Lys Tyr Trp Tyr Glu
 145 150 155 160
 40 Asn His Asn Ile Ser Ile Thr Asn Ala Thr Val Glu Asp Ser Gly Thr
 165 170 175
 45 Tyr Tyr Cys Thr Gly Lys Val Trp Gln Leu Asp Tyr Glu Ser Glu Pro
 180 185 190
 50 Leu Asn Ile Thr Val Ile Lys Ala Pro Arg Glu Lys Tyr Trp Leu Gln
 195 200 205
 55 Phe Phe Ile Pro Leu Leu Val Val Ile Leu Phe Ala Val Asp Thr Gly
 210 215 220
 60 Leu Phe Ile Ser Thr Gln Gln Gln Val Thr Phe Leu Leu Lys Ile Lys
 225 230 235 240
 65 Arg Thr Arg Lys Gly Phe Arg Leu Leu Asn Pro His Pro Lys Pro Asn
 245 250 255
 70 Pro Lys Asn Asn
 260
 75 <210> 11
 80 <211> 584
 85 <212> DNA
 90 <213> Homo sapiens

EP 1 006 183 A1

<220>
 <221> CDS
 <222> (12)..(569)
 5
 <220>
 <221> mat_peptide
 <222> (15)..(569)
 10
 <400> 11
 aatagaattc c atg ggg aca cct gca gct ccc cca aag gct gtg ctg aaa 50
 Met Gly Thr Pro Ala Ala Pro Pro Lys Ala Val Leu Lys
 -1 1 5 10
 15
 ctc gag ccc cag tgg atc aac gtg ctc cag gag gac tct gtg act ctg 98
 Leu Glu Pro Gln Trp Ile Asn Val Leu Gln Glu Asp Ser Val Thr Leu
 15 20 25
 20
 aca tgc cgg ggg act cac agc cct gag agc gac tcc att cag tgg ttc 146
 Thr Cys Arg Gly Thr His Ser Pro Glu Ser Asp Ser Ile Gln Trp Phe
 30 35 40
 25
 cac aat ggg aat ctc att ccc acc cac acg cag ccc agc tac agg ttc 194
 His Asn Gly Asn Leu Ile Pro Thr His Thr Gln Pro Ser Tyr Arg Phe
 45 50 55 60
 30
 aag gcc aac aac aat gac agc ggg gag tac acg tgc cag act ggc cag 242
 Lys Ala Asn Asn Asn Asp Ser Gly Glu Tyr Thr Cys Gln Thr Gly Gln
 65 70 75
 35
 acc agc ctc agc gac cct gtg cat ctg aca gtg ctt tct gag tgg ctg 290
 Thr Ser Leu Ser Asp Pro Val His Leu Thr Val Leu Ser Glu Trp Leu
 80 85 90
 40
 gtg ctc cag acc cct cac ctg gag ttc cag gag gga gaa acc atc gtg 338
 Val Leu Gln Thr Pro His Leu Glu Phe Gln Glu Gly Glu Thr Ile Val
 95 100 105
 45
 ctg agg tgc cac agc tgg aag gac aag cct ctg gtc aag gtc aca ttc 386
 Leu Arg Cys His Ser Trp Lys Asp Lys Pro Leu Val Lys Val Thr Phe
 110 115 120
 50
 ttc cag aat gga aaa tcc aag aaa ttt tcc cgt tcg gat ccc aac ttc 434
 Phe Gln Asn Gly Lys Ser Lys Lys Phe Ser Arg Ser Asp Pro Asn Phe
 125 130 135 140
 55
 tcc atc cca caa gca aac cac agt cac agt ggt gat tac cat tgc aca 482
 Ser Ile Pro Gln Ala Asn His Ser His Ser Gly Asp Tyr His Cys Thr
 145 150 155

EP 1 006 183 A1

gga aac ata ggc tac acg ctg tac tca tcc aag cct gtg acc atc act 530
 Gly Asn Ile Gly Tyr Thr Leu Tyr Ser Ser Lys Pro Val Thr Ile Thr
 160 165 170

gtc caa gct ccc agc tct tca ccg atg ggg atc att tag gctgtcgaca 579
 Val Gln Ala Pro Ser Ser Ser Pro Met Gly Ile Ile
 175 180 185

ctggg 584

<210> 12
 <211> 185
 <212> PRT
 <213> Homo sapiens

<400> 12
 Met Gly Thr Pro Ala Ala Pro Pro Lys Ala Val Leu Lys Leu Glu Pro
 1 5 10 15

Gln Trp Ile Asn Val Leu Gln Glu Asp Ser Val Thr Leu Thr Cys Arg
 20 25 30

Gly Thr His Ser Pro Glu Ser Asp Ser Ile Gln Trp Phe His Asn Gly
 35 40 45

Asn Leu Ile Pro Thr His Thr Gln Pro Ser Tyr Arg Phe Lys Ala Asn
 50 55 60

Asn Asn Asp Ser Gly Glu Tyr Thr Cys Gln Thr Gly Gln Thr Ser Leu
 65 70 75 80

Ser Asp Pro Val His Leu Thr Val Leu Ser Glu Trp Leu Val Leu Gln
 85 90 95

Thr Pro His Leu Glu Phe Gln Glu Gly Glu Thr Ile Val Leu Arg Cys
 100 105 110

His Ser Trp Lys Asp Lys Pro Leu Val Lys Val Thr Phe Phe Gln Asn
 115 120 125

Gly Lys Ser Lys Lys Phe Ser Arg Ser Asp Pro Asn Phe Ser Ile Pro
 130 135 140

Gln Ala Asn His Ser His Ser Gly Asp Tyr His Cys Thr Gly Asn Ile
 145 150 155 160

EP 1 006 183 A1

Gly Tyr Thr Leu Tyr Ser Ser Lys Pro Val Thr Ile Thr Val Gln Ala
 165 170 175
 5
 Pro Ser Ser Ser Pro Met Gly Ile Ile
 180 185
 10
 <210> 13
 <211> 570
 <212> DNA
 <213> Homo sapiens
 15
 <220>
 <221> CDS
 <222> (10)..(561)
 20
 <220>
 <221> mat_peptide
 <222> (13)..(561)
 25
 <400> 13
 gatggccat atg gca gtc cct cag aaa cct aag gtc tcc ttg aac cct cca 51
 Met Ala Val Pro Gln Lys Pro Lys Val Ser Leu Asn Pro Pro
 -1 1 5 10
 30
 tgg aat aga ata ttt aaa gga gag aat gtg act ctt aca tgt aat ggg 99
 Trp Asn Arg Ile Phe Lys Gly Glu Asn Val Thr Leu Thr Cys Asn Gly
 15 20 25
 35
 aac aat ttc ttt gaa gtc agt tcc acc aaa tgg ttc cac aat ggc agc 147
 Asn Asn Phe Phe Glu Val Ser Ser Thr Lys Trp Phe His Asn Gly Ser
 30 35 40 45
 40
 ctt tca gaa gag aca aat tca agt ttg aat att gtg aat gcc aaa ttt 195
 Leu Ser Glu Glu Thr Asn Ser Ser Leu Asn Ile Val Asn Ala Lys Phe
 50 55 60
 45
 gaa gac agt gga gaa tac aaa tgt cag cac caa caa gtt aat gag agt 243
 Glu Asp Ser Gly Glu Tyr Lys Cys Gln His Gln Gln Val Asn Glu Ser
 65 70 75
 50
 gaa cct gtg tac ctg gaa gtc ttc agt gac tgg ctg ctc ctt cag gcc 291
 Glu Pro Val Tyr Leu Glu Val Phe Ser Asp Trp Leu Leu Leu Gln Ala
 80 85 90
 55
 tct gct gag gtg gtg atg gag ggc cag ccc ctc ttc ctc agg tgc cat 339
 Ser Ala Glu Val Val Met Glu Gly Gln Pro Leu Phe Leu Arg Cys His
 95 100 105

EP 1 006 183 A1

5 ggt tgg agg aac tgg gat gtg tac aag gtg atc tat tat aag gat ggt 387
Gly Trp Arg Asn Trp Asp Val Tyr Lys Val Ile Tyr Tyr Lys Asp Gly
110 115 120 125

10 gaa gct ctc aag tac tgg tat gag aac cac aac atc tcc att aca aat 435
Glu Ala Leu Lys Tyr Trp Tyr Glu Asn His Asn Ile Ser Ile Thr Asn
130 135 140

15 gcc aca gtb gaa gac agt gga acc tac tac tgt acg ggc aaa gtg tgg 483
Ala Thr Val Glu Asp Ser Gly Thr Tyr Tyr Cys Thr Gly Lys Val Trp
145 150 155

20 cag ctg gac tat gag tct gag ccc ctc aac att act gta ata aaa gct 531
Gln Leu Asp Tyr Glu Ser Glu Pro Leu Asn Ile Thr Val Ile Lys Ala
160 165 170

25 ccg cgt gag aag tac tgg cta caa ttt tag gatccattg 570
Pro Arg Glu Lys Tyr Trp Leu Gln Phe
175 180

30 <210> 14
<211> 183
<212> PRT
<213> Homo sapiens

35 Met Ala Val Pro Gln Lys Pro Lys Val Ser Leu Asn Pro Pro Trp Asn
1 5 10 15

40 Arg Ile Phe Lys Gly Glu Asn Val Thr Leu Thr Cys Asn Gly Asn Asn
20 25 30

45 Phe Phe Glu Val Ser Ser Thr Lys Trp Phe His Asn Gly Ser Leu Ser
35 40 45

50 Glu Glu Thr Asn Ser Ser Leu Asn Ile Val Asn Ala Lys Phe Glu Asp
50 55 60

55 Ser Gly Glu Tyr Lys Cys Gln His Gln Gln Val Asn Glu Ser Glu Pro
65 70 75 80

Val Tyr Leu Glu Val Phe Ser Asp Trp Leu Leu Leu Gln Ala Ser Ala
85 90 95

60 Glu Val Val Met Glu Gly Gln Pro Leu Phe Leu Arg Cys His Gly Trp
100 105 110

55

Arg Asn Trp Asp Val Tyr Lys Val Ile Tyr Tyr Lys Asp Gly Glu Ala
 115 120 125
 5
 Leu Lys Tyr Trp Tyr Glu Asn His Asn Ile Ser Ile Thr Asn Ala Thr
 130 135 140
 10
 Val Glu Asp Ser Gly Thr Tyr Tyr Cys Thr Gly Lys Val Trp Gln Leu
 145 150 155 160
 Asp Tyr Glu Ser Glu Pro Leu Asn Ile Thr Val Ile Lys Ala Pro Arg
 15 165 170 175
 Glu Lys Tyr Trp Leu Gln Phe
 180
 20

Claims

- 25 1. Recombinant soluble Fc receptor
characterized by the absence of transmembrane domains, signal peptide and glycosylation.
- 30 2. Recombinant Fc receptor according to claim 1,
wherein the receptor is a FcγR or a FcεR.
- 3 3. Recombinant Fc receptor according to claim 1 or 2,
wherein the receptor is a FcγRIIb.
- 35 4. Recombinant Fc receptor according to any one of claims 1 to 3,
wherein the receptor is of human origin.
- 5 5. Recombinant Fc receptor according to any one of claims 1 to 4,
wherein it contains the amino acids as shown in SEQ ID NO: 1 or SEQ ID NO:2.
- 40 6. Recombinant nucleic acid containing a sequence encoding a recombinant Fc receptor according to any one of
claims 1 to 5.
- 7 7. Recombinant nucleic acid according to claim 6,
wherein it contains a sequence as shown in SEQ ID NO:3 or SEQ ID NO:4.
- 45 8. Recombinant nucleic acid according to claim 6 or 7,
wherein it additionally contains expression control sequences operably linked to the sequence encoding the recom-
binant Fc receptor.
- 50 9. Recombinant nucleic acid according to any one of claims 6 to 8,
wherein it is contained on a prokaryotic expression vector, preferably a pET vector.
- 10 10. Host cell characterized by the presence of a recombinant nucleic acid according to any one of claims 6 to 8.
- 55 11. Host cell according to claim 10,
wherein it is a prokaryotic host cell, preferably an E. coli cell.

12. Process for the determination of the amount of antibodies of a certain type in the blood, plasma or serum of a patient, characterized by the use of a recombinant soluble Fc receptor according to any one of claims 1 to 5 in an immunoassay and determination of the presence of FcR-antibody complexes.
- 5 13. Process according to claim 12, wherein the immunoassay is an ELISA and preferably a sandwich assay.
14. Process according to claim 12 or 13, wherein the antibodies to be determined are IgE antibodies and the recombinant soluble receptor is a FcεR.
- 10 15. Process according to claim 14 for the determination of a predisposition or manifestation of an allergy.
16. Process according to claim 12 or 13, wherein the antibodies to be determined are IgG antibodies and the recombinant soluble receptor is a FcγR.
- 15 17. Process for the determination of the immune status of patients with chronic diseases of the immune system, wherein a Fc receptor according to any one of claims 1 to 5 is used in a competitive immunoassay and the amount of the corresponding sFcRs in the blood, plasma or serum of a patient is determined.
18. Process according to claim 17, wherein the chronic disease is AIDS, SLE, MM or rheumatoid arthritis.
- 20 19. Use of a recombinant soluble Fc receptor according to any one of claims 1 to 5 for the screening of substances in view of their ability to act as inhibitors of the recognition and binding of antibodies to the respective cellular receptors.
- 25 20. Use according to claim 19, wherein recombinant soluble FcγRs are used and recognition and binding of IgG antibodies is of interest.
21. Pharmaceutical composition containing as active agent a recombinant soluble FcR according to any one of claims 1 to 5.
- 30 22. Pharmaceutical composition according to claim 21 for use in the treatment or prevention of autoimmune diseases, allergies or tumor diseases.
- 35 23. Pharmaceutical composition according to claim 21 or 22 for use in the treatment of AIDS, rheumatoid arthritis or multiple myeloma, containing a recombinant soluble FcγR preferably having the amino acid sequence as shown in SEQ ID NO:1.
24. Use of a crystalline preparation of a recombinant soluble Fc receptor according to any one of claims 1 to 5 for the generation of crystal structure data of Fc receptors.
- 40 25. Use of crystal structure data obtained by the use according to claim 24 for the identification and preparation of Fc receptor inhibitors.
26. Use of crystal structure data obtained by the use according to claim 24 for the identification and preparation of new antibody receptors.
- 45 27. Use according to any one of claims 24 to 26 in a computer-aided modelling program.
28. FcR inhibitor characterized in that it has a three-dimensional structure which is complementary to the recombinant soluble FcR according to any one of claims 1 to 5.
- 50 29. Pharmaceutical composition containing as active agent a FcR inhibitor according to claim 28.
30. Pharmaceutical composition according to claim 29 for use in the treatment or prevention of diseases which are due to overreactions or faulty reactions of the immune system.
- 55 31. Pharmaceutical composition according to claim 29 or 30 for the treatment or prevention of allergies, autoimmune diseases or an anaphylactic shock.

32. Fc receptor according to claims 1-5, bound to a solid phase.

33. Fc receptor according to claim 32, wherein the solid phase is a chromatography carrier material.

5 34. Use of a chromatography carrier material according to claim 33 for the adsorption of immunoglobulins from the blood, plasma or serum of a patient or from culture supernatants of immunoglobulin producing cells.

35. Use according to claim 34 for the enrichment of antibodies from a patient's blood, serum or plasma or from culture supernatants of immunoglobulin producing cells for the conduction of further tests.

10

15

20

25

30

35

40

45

50

55




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5A




FIG. 5B




FIG. 6




FIG. 7

Alignment of the amino acid sequence of the extracellular parts of FcγR and FcεR1a
(without signal sequence and transmembrane region)

1	50	100							
FcγRIIa	SQAAAAPPKAV LKLEPPWINV LQEDSVTLTC QGASPESDS IQWFHNGNLI PTHQTPSYRP KANNDSGEY TCQTGQTSL	DPVHLTVLSE WLVLTQPHLE							
FcγRIIb	GTPAAPPKAV LKLEPQWINV LQEDSVTLTC RGTSPESDS IQWFHNGNLI PTHQTPSYRF KANNDSGEY TCQTGQTSL	DPVHLTVLSE WLVLTQPHLE							
FcγRIII	MRTEDLPKAV VFLEPQWYSV LEKDSVTLKC QGAYSPEDNS TQWFHNSLI SSQASSYFID AATVNDSGEY RCQTNLSTLS	DPVQLEVHIG WLLIQAPRAW							
FcγRIAV ISLQPPWVSV FQESTVTLHC EVLHLPGSS TQWFLNGTAT QTSTPSYRIT SASVNDSGEY RCQRLSGRS	DPIQLEIHRG WLLIQVESRV							
FcεRIa	GVLAVPQKPK VSLNPPWNR	FKGENVTLTC NGNNEFEVSS TKWFHNGSL	ETNSSLNIV NAKFEDSGEY KCQHQQVNES EPVYLEVFS	WLLIQASAEV					
101	150	200							
FcγRIIa	FOEGETIMLR CHSWKDKPLV KVTFPQNGKS QKFSRLDPTF	SIPQANHSHS GDVHCTGNIG YTLFSSKPV	ITVQVPSMGS SSPMGII					
FcγRIIb	FOEGETIVLR CHSWKDKPLV KVTFPQNGKS KKFSRSDPNF	SIPQANHSHS GDVHCTGNIG YTLFSSKPV	ITVQAP...S SSPMGII					
FcγRIII	FKBEDPIHLR CHSWKNTALH KVTYLQNGKD RKYTHNSDF	HIPKATLKDS GSYFCRGLVG SKVSSSETVN	ITITQGLAVS TISSFPP					
FcγRI	FTEGEPLALR CHAWKDKLVY NVLYVRNGKA FKFFHNSNL	TILKTNISHN GTHCSG.MG KHRYSAGIS	VTVKELFPAP VLNASVTSPL	LEGNLVTLS					
FcεRIa	VMEGQPLFLR CHGRNWDVY KVIYKOGEEA LKYWYENHNI	SITNATVEDS GTTYCTGKVN	QLDYESEPLN	ITVIKAPREK YWLQ.....				
201	250	277							
FcγRIIa				
FcγRIIb				
FcγRIII				
FcγRI	ETKILLQRP	LQLYFSFYM	SKTLGRNTS	SEYQILTARR	EDSGLYWCEA	ATEDGNVLKR	SPELEQLVG	LQLPTPV
FcεRIa

FIG. 8



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	EP 0 614 978 A (ROUSSEL-UCLAF) 14 September 1994 * page 3, line 1 - page 7, line 33; examples *	1-3, 6, 8-11, 21-23	C12N15/12 C07K14/705 C12N1/21 C12N15/70 G01N33/53 G01N33/68
X	US 5 623 053 A (LOUIS N. GASTINEL ET AL.) 22 April 1997 * column 5, line 60 - column 8, line 67; examples *	1, 4, 6, 8-13, 24-27	A61K38/17 C07K17/00
X	WO 96 40199 A (UNIVERSITY OF PENNSYLVANIA) 19 December 1996 * page 22, line 24 - page 24, line 14; claims 13-16, 36-39; examples I, II *	1, 2, 4, 6, 8-16, 19-22, 32-35	
X	WO 95 09002 A (UNIVERSITY OF PENNSYLVANIA) 6 April 1995 * page 15, line 4 - page 16, line 22; claims 28-31; examples I, II *	1, 2, 4, 6, 8-13, 16, 21, 32, 33	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07K C12N G01N A61K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely:</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>28-31</p> <p>Reason for the limitation of the search:</p> <p>Subject-matter directed to inhibitors has not been searched due to the lack of adequate technical description thereof in the application.</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		6 May 1999	Montero Lopez, B
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.92 (P04C07)



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 98 12 2969

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	EP 0 791 653 A (SCHERING BIOTECH CORPORATION) 27 August 1997 * column 5, line 56 - column 10, line 1; example III *	1,2,4,6, 8-11,21, 22	
X	JÉRÔME GALON ET AL.: "Ligands and biological activities of soluble Fcγ receptors" IMMUNOLOGY LETTERS, vol. 44, January 1995, pages 175-181, XP000574018 * page 175, right-hand column, last paragraph - page 177, left-hand column, paragraph 1 * * page 179, right-hand column, paragraph 3 - page 180, right-hand column, paragraph 3 *	1-4,6, 8-13,15, 16,19-22	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
X	WO 92 01049 A (THE GENERAL HOSPITAL CORPORATION) 23 January 1992 * page 10, line 18 - page 11, line 19 * * page 13, line 11 - page 14, line 4; example X; table 2 *	5-11	
E	WO 99 05271 A (GOULD, HANNAH ET AL.) 4 February 1999 * page 4, line 6 - line 19 * * page 5, line 22 - page 6, line 2 * * page 7, line 22 - page 8, line 23 * * page 10, line 31 - page 11, line 31 *	1,2,4,6, 8-15,21, 22,32	

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 12 2969

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-05-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 614978 A	14-09-1994	FR 2702481 A	16-09-1994
		CA 2118558 A	10-09-1994
		JP 7149800 A	13-06-1995
US 5623053 A	22-04-1997	NONE	
WO 9640199 A	19-12-1996	AU 6386996 A	30-12-1996
		CA 2223402 A	19-12-1996
		EP 0831875 A	01-04-1998
		US 5858981 A	12-01-1999
WO 9509002 A	06-04-1995	AU 701919 B	11-02-1999
		AU 7924494 A	18-04-1995
		CA 2172392 A	06-04-1995
		EP 0723455 A	31-07-1996
		JP 9506072 T	17-06-1997
		US 5858981 A	12-01-1999
EP 791653 A	27-08-1997	AU 3740089 A	12-12-1989
		EP 0343950 A	29-11-1989
		EP 0449828 A	09-10-1991
		JP 2657221 B	24-09-1997
		JP 3501484 T	04-04-1991
		KR 149012 B	17-08-1998
		WO 8911490 A	30-11-1989
		US 5767077 A	16-06-1998
		CA 2000355 A	19-07-1990
WO 9201049 A	23-01-1992	AU 658370 B	13-04-1995
		AU 8528691 A	04-02-1992
		CA 2087272 A	14-01-1992
		EP 0551301 A	21-07-1993
		JP 6504186 T	19-05-1994
		PT 98326 A, B	30-06-1992
		US 5506126 A	09-04-1996
		US 5849898 A	15-12-1998
		US 5830731 A	03-11-1998
WO 9905271 A	04-02-1999	NONE	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.